

1-1-2005

# Clinical lactation practice: 20 years of evidence

Karen Wambach

Suzanne H. Campbell

*Fairfield University*, [suzanne.campbell@nursing.ubc.ca](mailto:suzanne.campbell@nursing.ubc.ca)

Sara Gill

Joan Dodgson

Titi Abiona

*See next page for additional authors*

This is the pre-peer reviewed version of the following article: "Clinical lactation practice: 20 years of evidence" which has been published in final form in the *Journal of Human Lactation* ( 21, 245-258)  
DOI: 10.1177/0890334405279001 – Copyright 2005 Sage Publications.

---

## Repository Citation

Wambach, Karen; Campbell, Suzanne H.; Gill, Sara; Dodgson, Joan; Abiona, Titi; and Heinig, M. Jane, "Clinical lactation practice: 20 years of evidence" (2005). *Nursing Faculty Publications*. Paper 2.  
<http://digitalcommons.fairfield.edu/nursing-facultypubs/2>

## Published Citation

Wambach, K., Campbell, S. H., Gill, S., Dodgson, J., Abiona, T., Heinig, M. J. (2005). Clinical lactation practice: 20 years of evidence. *Journal of Human Lactation*, 21, 245-258.

This Article is brought to you for free and open access by the School of Nursing at DigitalCommons@Fairfield. It has been accepted for inclusion in Nursing Faculty Publications by an authorized administrator of DigitalCommons@Fairfield. For more information, please contact [digitalcommons@fairfield.edu](mailto:digitalcommons@fairfield.edu).

---

**Authors**

Karen Wambach, Suzanne H. Campbell, Sara Gill, Joan Dodgson, Titi Abiona, and M. Jane Heinig

## Clinical Lactation Practice: 20 Years of Evidence

Karen Wambach, Suzanne Campbell, Sara Gill, Joan Dodgson, Titi Abiona , and M. Jane Heinig

International Board Certified Lactation Consultants come from many backgrounds including nursing, midwifery, medicine, medical and nursing education, public health, occupational therapy, anthropology, dietetics and nutrition, speech therapy and pathology, physical therapy, and social work. As practitioners devoted to promoting and supporting breastfeeding among childbearing women, lactation consultants depend on a vast multidisciplinary knowledge base to support their practices, with evidence-based knowledge considered the gold standard.<sup>1</sup>

Over the past twenty years there has been an evolution of methods used in breastfeeding related research. This evolution has been a product of several factors including: 1) an increased knowledge base which results in the increased use of controlled experimental designs;<sup>2,3</sup> 2) increased attention to definitions of infant feeding practices<sup>4</sup> and outcomes; 3) increased opportunities for funding (affecting methods, sample size, and sophistication); and 4) multi-disciplinary collaboration. As the evidence for breastfeeding/human milk's health benefits has grown, so have the efforts to improve breastfeeding rates and duration. In this review, we will summarize the major findings of research investigations conducted within selected areas (health outcomes, factors associated with initiation and duration of breastfeeding, common problems, special populations, and infant feeding practices in developing countries) during the last 20 years.

For this review, the English language scientific literature (CINAHL, MedLine, Cochrane Database of Systematic Reviews) was searched for relevant articles published since 1985.

### Health Consequences of Infant Feeding Practices for Mothers and Infants

Since 1985, there have been significant advances in our understanding of health outcomes associated with breastfeeding. Human milk confers protection to the infant against infectious agents by a unique blend of bioactive constituents that work alone or in combination to develop and enhance the infant's immature immune system.<sup>5</sup> Research into these factors, their functions, and sites of action continues today. Human milk contains soluble and cellular factors from the maternal immune system, antimicrobial substances and developmental factors that assist the infant to resist infection.<sup>6</sup> Examples of some of these constituents may be found in Table 1. The protection conferred by human milk is unique for each mother and infant because the exposure of the mother to environmental antigens results in the transfer of immune factors to the milk that are specific to these antigens.<sup>7,8</sup>

Epidemiological studies support the association between breastfeeding and reduced infant morbidity and mortality in both developing and industrialized nations, particularly against otitis media and gastrointestinal and lower respiratory illnesses.<sup>5,9-12</sup> In early studies, the ability of researchers to find differences between breastfed and artificially fed populations was challenged by the classification of infants as "breastfed" no matter how much or little breast milk they had received. In those early studies, infants who breastfed only once or twice often were grouped with infants who were exclusively breastfed for several months.<sup>10</sup> In 1990, Labbok and Krasovec published a classification

schema of infant feeding practices to be used by researchers examining the impact of breastfeeding on both mothers and infants.<sup>13</sup> A modified scheme for classifying infant feeding was later adopted and disseminated by the World Health Organization.<sup>14</sup> It is now known that examining not only if an infant was breastfed but how (in terms of duration and exclusivity) is essential to our understanding of the impact of breastfeeding on human health. The evidence indicates that the greatest protection conferred by breastfeeding against infectious illness occurs when breastfeeding is exclusive and continues for at least 3 months.<sup>5</sup> Based on extensive evidence related to the advantages of breastfeeding for infant health and development, it was concluded by many health organizations including the WHO<sup>14</sup> and the American Academy of Pediatrics,<sup>12</sup> that exclusive breastfeeding should continue for the first 6 months of life. Continued breastfeeding, with complementary foods is encouraged for at least 12 months and as long as is desired.

Human milk is also associated with improved health in premature infants,<sup>15</sup> and higher scores on tests of cognitive function than those of artificially-fed preterm infants.<sup>16,17</sup> In a meta-analysis of studies among preterm low-birth weight infants, those who received donor milk were 3 times less likely to develop necrotizing enterocolitis than those who received formula.<sup>18</sup> Further work is needed to examine the effects of breast milk feeding among very-low-birth weight infants.<sup>19</sup>

Evidence indicates that breastfeeding may be protective against chronic disease, such as some childhood cancers,<sup>20</sup> atopic disease,<sup>21,22</sup> diabetes,<sup>20</sup> and childhood obesity.<sup>23</sup> Breastfeeding is also associated with higher cognitive development scores

among term infants as compared to artificially-fed infants among many but not all studies.<sup>24-26</sup>

For mothers, breastfeeding results in both short and long term physiological benefits. Initially, breastfeeding causes the uterus to contract, reducing the mothers' risk for postpartum hemorrhage and hastening her recovery from labor.<sup>27</sup> With increasing duration of breastfeeding, mothers gain protection from postpartum weight retention,<sup>28</sup> breast cancer,<sup>27-29</sup> and in some studies, ovarian cancer.<sup>30</sup> Among populations that do not use hormonal contraceptives, breastfeeding serves as an important method of contraception.<sup>27</sup>

Despite all the knowledge that we have gained in the past 20 years about the impact of breastfeeding on the infant and mother, many families still choose not to breastfeed. The identification of factors associated with breastfeeding initiation and duration has been another important area of research.

### **Getting Mothers Started: Factors and Interventions Associated with Breastfeeding Initiation**

The factors associated with initiation of breastfeeding have been widely researched using various quantitative (e.g., large-scale national surveys, regional and local observational studies) and qualitative (e.g., descriptive experiences of mothers, healthcare providers, etc.) methods. Maternal experiences, personal, socio-cultural and economic characteristics and types of support available (e.g., informal, formal and environmental) have been explored. Many researchers have agreed upon maternal characteristics (i.e., mothers who are older, better educated, more affluent, and married) and personal attributes (e.g., positive attitudes and beliefs toward breastfeeding,

persistence, confidence) most often associated with breastfeeding initiation.<sup>31-35</sup>

Negative breastfeeding experience is associated with choosing not to breastfeed subsequent children.<sup>36</sup>

Mothers' need to return to work during the early postpartum is a barrier to breastfeeding<sup>37-41</sup> and may particularly affect women working in conditions where pumping is not a feasible option.<sup>42</sup> However, in some non-Western countries more affluent women choose not to breastfeed as a sign of their wealth.<sup>43, 44</sup>

Researchers have identified the lack of informal (e.g., family, social network), formal (e.g., healthcare providers) and environmental (e.g., places to breastfeed while out of the house, public media) support as negatively affecting breastfeeding initiation.<sup>41,45-48</sup>

Interventions intended to increase breastfeeding initiation have yielded mixed results. National media campaigns have been used internationally to improve public perceptions about breastfeeding.<sup>49</sup> Prenatal education and peer support also are interventions widely associated with increasing breastfeeding initiation rates even among groups who do not typically choose to breastfeed.<sup>33,41,50-53</sup> However, these programs have not been consistently effective.<sup>33,54-55</sup> Socio-economic disparities may affect a woman's access to prenatal breastfeeding education.<sup>56</sup> While still lower than the general population, US women who have traditionally breastfed at lower rates (i.e., teenaged, African American, participants in WIC or Medicaid, those who obtain late prenatal care or have an infant in NICU) are initiating breastfeeding in greater numbers<sup>57,58</sup> perhaps due to increased public and private breastfeeding promotion programs.<sup>58</sup> Similar results have been reported in countries where hospital practices have actively incorporated the Baby-Friendly Hospital Initiative.<sup>41</sup> Our knowledge about what is most effective in

promoting breastfeeding initiation would be improved if more rigorous research was done on various interventions.<sup>9</sup>

### **Increasing Duration and Exclusivity of Breastfeeding: Factors and Interventions**

Surveillance of national rates of exclusive breastfeeding is included in the WHO Global Data Bank on Breastfeeding.<sup>59</sup> An issue in conducting research measuring breastfeeding exclusivity is the inaccuracy of maternal recall.<sup>60</sup> Factors associated with duration of exclusive breastfeeding are similar in many countries. Among US women, duration of exclusive breastfeeding is associated with primiparity, non-Hispanic African American ethnicity, less than 20 years of age, lower income, lower education and type of employment.<sup>61</sup> A population-based survey in Quebec, Canada found maternal age to be the predominate determinant of exclusivity.<sup>62</sup> An increased risk of early weaning occurs with maternal smoking according to a meta-analysis of the effect of smoking on breastfeeding.<sup>63</sup>

Some researchers have examined the factors influencing breastfeeding duration by using psychosocial theories (e.g., theory of planned behavior, attachment, self-efficacy) that hypothesize the relationships between internal psychological and external socio-environmental influences on women's infant feeding behaviors.<sup>32,34, 64-66</sup> They report increased breastfeeding duration is associated with positive intention, attitudes, beliefs, and increased social support.

Health education programs regarding the benefits of breastfeeding have been effective in increasing breastfeeding initiation rates, but much less effective in maintaining duration and exclusivity.<sup>33,45</sup> Supportive early postpartum experiences, particularly while in the hospital, have been associated with increased duration and length



of exclusivity.<sup>45</sup> Breastfeeding support also has been cited as a post-discharge intervention effective in increasing both duration and exclusivity, particularly when obtained from a health care provider.<sup>67</sup> Lay breastfeeding support has not been as effective,<sup>67</sup> though support from teams of professional health care providers and lay/peer counselors have been successful in increasing breastfeeding duration.<sup>68,69</sup> Lack of skilled professional support also is a factor in early weaning and decreased exclusivity.<sup>70,71</sup> Improving our understanding of how best to facilitate exclusive breastfeeding is an essential area for future research.

### **The Science of Breastfeeding Support: Advances in Our Knowledge of Mammary Physiology**

While many of the major structures of the mammary gland were described more than 50 years ago, advances in technology have led to a greater understanding of mammary function at the cellular and molecular levels.<sup>72</sup> Extensive work has been ongoing to gain greater understanding of the anatomy and variation in the functioning mammary gland as well as the nature of the functions of the gland, control over those functions, and the interaction of the gland with the rest of the body.<sup>72-74</sup>

Much of what we know about the onset of milk production and control of milk volume is based on research that was conducted in the last 20 to 25 years. Hartmann and others have divided the onset of milk production (lactogenesis) into 2 parts, stages I and II.<sup>75,76</sup> During stage I which occurs during pregnancy, mammary gland structures are differentiated into functional cells capable of secretion, and plasma concentrations of lactose and  $\alpha$ -lactalbumin increase. Glandular secretion is inhibited by progesterone and to a lesser extent, estrogen. Though high levels of prolactin are needed for lactogenesis,

the surge in prolactin that occurs after parturition is not likely the trigger for milk onset.<sup>77</sup> Colostrum is secreted during pregnancy and in the early postpartum period. Colostrum is higher in protein, minerals and immune factors, and lower in fat as compared to mature milk.<sup>78</sup>

During stage II, there is an onset of copious milk production that occurs in the first 4 days postpartum, as progesterone levels drop 10 fold. There are rapid changes in milk volume and composition and the transition from colostrum to mature milk occurs on a continuum. Mammary structures continue to change resulting in reduction in permeability of the paracellular pathway between epithelial cells (junctions) and increased secretion of all milk components. As lactation progresses, prolactin levels fall and are not correlated with milk volume. A complex system of hormonal interactions is needed to maintain lactation<sup>74</sup> and autocrine systems adjust milk volume in response to milk removal by the infant.<sup>76</sup>

Milk secretion and plasma changes occur whether or not women choose to breastfeed, suckling and milk removal are not necessary for lactogenesis II to occur. However, if milk removal does not occur around day 3 postpartum, milk secretion will typically decline. Time of first breastfeeding and frequency of breastfeeding on day 2 are positively correlated with milk volume on day 5. Formula feeding prior to lactogenesis is associated with significant delay onset of stage II, suggesting that milk removal is important in increasing the efficiency of milk secretion.<sup>79</sup> Other factors associated with delay of milk onset include primiparity, maternal body mass index greater than 27 kg/m<sup>2</sup>, placental retention (progesterone), cesarean section delivery, and elevated cortisol (stress during parturition).<sup>80,81</sup>

Delay of lactogenesis is only one of several challenges commonly faced by lactating women. An understanding of the nature of these challenges is needed to enhance the support of the breastfeeding dyad.

### **What Are The Most Common Challenges For Breastfeeding Mothers?**

A number of maternal and infant based concerns or problems may be encountered during breastfeeding and adversely affect breastfeeding success. Maternal based concerns include sore nipples, engorgement, mastitis, and perceived or actual insufficient milk supply. Due to space constraints, infant based concerns such as suck disorders and hyperbilirubinemia will not be addressed in this paper.

**Sore nipples** are one of the most common reasons that women discontinue breastfeeding in the early postpartum period.<sup>82-83</sup> Although some research has suggested that nipple changes in the first weeks of breastfeeding are common and normal,<sup>84, 85</sup> more often, faulty breastfeeding techniques, including improper positioning, latch-on, and artificial nipple use, are implicated in nipple trauma and pain.<sup>86-88</sup> Staphylococcus infection also is associated with sore, cracked nipples.<sup>89</sup> Oral anomalies, including ankyloglossia, and suckling disorders in the infant have also been implicated in the development of sore nipples.<sup>90,91</sup>

Clinical trials conducted internationally to investigate the effectiveness of various treatment methods for sore nipples such as lanolin, hydrogel dressings, tea bag and water compresses, expressed breast milk, air-drying, breast shells, and education have had mixed results. Results of trials testing lanolin nipple ointments, alone, or in comparison to other preparations have demonstrated neutral results,<sup>92,93</sup> negative results;<sup>94</sup> and positive results.<sup>95</sup> Breast infections have been associated with both lanolin<sup>94</sup> and hydrogel

dressings.<sup>95</sup> Lavergne<sup>96</sup> found that tea bag and water compresses were more effective than no treatment, and there was no statistically significant difference between the two types of compresses. Future research should include multiple methods of treatment, in addition to, and separately from education interventions.

**Engorgement** is also a problem that breastfeeding mothers experience. Descriptive research by Humenick and colleagues<sup>97,98</sup> described patterns of engorgement and identified correlates of significance (initiation of feeding, frequency of feedings, feeding duration, rate of milk maturation, and supplementation). Clinical trials to test various methods of engorgement treatment have been conducted. A meta-analysis by the Cochrane Review group reviewed eight trials, involving 424 women.<sup>99</sup> No overall benefit was identified in three different studies that used cabbage leaves or cabbage leaf extracts. Ultrasound treatment and placebo were equally effective. Use of Danzen (an anti-inflammatory agent) significantly improved the total engorgement symptoms when compared to placebo as did bromelain/trypsin complex. Oxytocin and cold packs had no apparent effect on engorgement symptoms. Additional research is needed in prevention and treatment of engorgement.

**Lactation mastitis** is a common inflammatory condition of the breast with prospective descriptive studies reporting incidences of 9.5% to 23.7% over periods of 3, 6, and 12 months postpartum.<sup>100-103</sup> Risk factors identified in prospective research include a history of mastitis with a previous child, cracked and sore nipples, and plugged ducts.<sup>101,103-105</sup> Most commonly, *Staphylococcus aureus* is the causative organism.<sup>89,105-107</sup> Often occurring early in the puerperium, the onset is generally abrupt, and causes significant discomfort and disruption for the breastfeeding mother;<sup>108-109</sup> sometimes

leading to weaning.<sup>100,103,105,109</sup> Mastitis recurrence is common.<sup>100,103,109</sup> Mothers report the most common and effective means of self-care include breast massage, frequent feeding from the affected breast, bed rest, analgesics/anti-inflammatory agents, hot packs, and compliant antibiotic use.<sup>108,109</sup>

**Insufficient Milk** is consistently reported as the most common reason for weaning. The problem is one of complexity and crosses international, cultural, and socioeconomic lines.<sup>110-113</sup> Early work in describing the phenomenon occurred concurrently and prior to the establishment of ILCA (and this literature review boundary). However, research related to biological and behavioral predictors of milk volume has been ongoing with notable work conducted in the 1990s.<sup>114-122</sup> More recently, psychological factors such as maternal confidence and self-efficacy have been investigated.<sup>64, 122</sup> Efforts to systematically reduce the incidence of insufficient milk have been employed through efforts such as ILCA's *Evidence Based Guidelines For the First 14 days of Life*.<sup>123</sup> However, no clinical trial with the primary aim of preventing or treating insufficient milk was found.

## **How Do We Best Promote and Support Breastfeeding In Special And Vulnerable Populations?**

### **Preterm / Very Low Birth Weight (VLBW) Infants**

Research-based knowledge in the area of breastfeeding and the premature/very low birth weight infant has increased phenomenally over the past twenty years. The preterm infant has very special needs and research continues to produce evidence of the superiority of human milk for the preterm infant in comparison to commercially prepared preterm formula in the areas of host defense, gastrointestinal function, nutrition, and neurodevelopment.<sup>124</sup> Although mothers of preterm infants face numerous challenges in

providing their own milk for their infants.<sup>125, 126</sup> they report that their efforts are outweighed by the rewards.<sup>127,128</sup>

Seminal early work in the 1980's provided much needed evidence to demonstrate the physiologic ability of preterm infants to suckle at the breast without physiologic compromise in comparison to bottle-feeding.<sup>129-131</sup> Concurrently and thereafter, research relative to milk expression via pumping added to the knowledge base for NICU lactation support teams.<sup>132-134</sup> Because of preterm infants' special nutrition and caloric needs, the science of lacto-engineering was used to enhance the milk fat content in human milk.<sup>135</sup> Efforts to improve milk transfer using special devices such as nipple shields also were examined<sup>136</sup> as were optimal support strategies for mothers of non-nursing preterm infants.<sup>137</sup> Research on the use of kangaroo-care for enhancing breastfeeding in the preterm infant has also grown in recent years.<sup>138-140</sup> Additionally, information regarding the behavioral responses of the preterm infant to breastfeeding has enhanced the holistic lactation care of the preterm infant and mother.<sup>141-144</sup>

### **Adolescent Mothers**

Typically, breastfeeding initiation rates and duration among adolescent mothers are lower than among adult women. Although health care professionals and others may question teen mothers' maturity and commitment to breastfeeding,<sup>145</sup> teenage mothers are physiologically able to lactate as well as their adult counterparts and without ill effects to their own physical growth and development.<sup>146-149</sup> The benefits of human milk are especially important in this vulnerable population because infants of teenage mothers are hospitalized in the first year of life more often than infants of older mothers, often for infections such as gastroenteritis.<sup>150,151</sup>

Research indicates that teenagers' attitudinal, social, ethnic/racial, perceived control, and commitment factors are influential in choosing breast or bottle-feeding.<sup>152-159</sup> Teens breastfeeding experiences and challenges are often similar in nature to those of adult women including physical breast discomforts,<sup>152,160-162</sup> concerns about insufficient milk<sup>147,160-162</sup> and fatigue.<sup>161</sup> Breastfeeding problems thought to be more specific to adolescent mothers include modesty issues, maintenance of breastfeeding after return to school, trouble expressing breast milk, and maternal sleep interruptions.<sup>147,152,160-163</sup> Teens also report the positive aspects of breastfeeding, including the closeness of the nursing relationship or bonding,<sup>163</sup> and ease.<sup>160</sup> Finally, reasons for weaning identified in the teen mother include: infant related reasons such as improper latching to the breast;<sup>163</sup> a demanding, unsettled, or rejecting infant;<sup>147, 162</sup> insufficient milk supply and baby liking the bottle better;<sup>152,162</sup> and social influences such as inaccurate advice, relatives' and doctors' advice to quit, embarrassment, and work or school barriers.<sup>152,160,162</sup>

Very few clinical trials focusing on promotion and support of breastfeeding in teens have been conducted. Using a combination of education and professional and peer counseling, Volpe and Bear<sup>164</sup> demonstrated an increase in breastfeeding initiation to 65% ( $p < .001$ ) from the previous rate of 14% (historical control group). Therefore, additional clinical trials using developmentally appropriate interventions are needed to enhance breastfeeding initiation and duration in teenage mothers.

### **The effect of ethnicity and low-income on breastfeeding**

Researchers have emphasized the important influence of cultural influences on breastfeeding practices.<sup>165-171</sup> Cultural understanding can be very important in promotion of breastfeeding initiation and is essential to providing services.<sup>170</sup> Unfortunately, few

reports have been published describing culturally relevant modes of information and service.<sup>41,172,173</sup>

According to the 2001 U.S. National Immunization Survey, non-Hispanic blacks had the lowest rates of breastfeeding initiation and continuation in the US. Exclusive breastfeeding rates were also lowest in this group.<sup>174</sup> African American women stated that they did not breastfeed for the following reasons: loss of breast shape; insufficient milk supply; loss of freedom; perception that partner doesn't like it; and public embarrassment.<sup>175</sup> The prenatal period may be the most important time to impress black women with the benefits of breastfeeding.<sup>176</sup>

Breastfeeding rates are also low among Hispanic women in the US.<sup>177</sup> The more acculturated Hispanic women are to the United States, the less likely they are to breastfeed.<sup>172,177,178</sup> Age, degree of acculturation, and marital status were significantly associated with breastfeeding initiation, while education and income were not associated with initiation among a group of Hispanic women on the Texas/Mexico border.<sup>178</sup>

Hispanic women in the US are aware of the benefits of breastfeeding, but knowledge of benefits to the mother and baby may not overcome perceived problems such as pain, embarrassment and inconvenience.<sup>179, 180</sup> In addition these low-income mothers identify several concerns related to breastfeeding including lack of confidence, loss of freedom, lifestyle restrictions, and lack of support from family and friends.<sup>181</sup>

Research identifies the women least likely to breastfeed as young, low-income members of ethnic minorities and those who lack support for breastfeeding.<sup>45,182</sup> Women at risk of premature cessation decide to breastfeed later in their pregnancy, demonstrate a negative attitude toward breastfeeding, positive attitudes about bottle feeding, and have a



low confidence in their ability to breastfeed.<sup>45, 183</sup> Early weaning was also associated with an intention to breastfeed a shorter period of time, lower breastfeeding knowledge scores, perceived insufficient milk supply, and plans to work outside the home.<sup>183</sup> Other studies have examined who influences low-income, ethnically diverse women's decisions regarding infant feeding choices, such as mothers and WIC personnel.<sup>152,184</sup> Studies indicate that additional support measures such as peer and paternal support may increase breastfeeding initiation and continuation among low-income diverse populations.<sup>185-186</sup>

### **Health professional support of breastfeeding women**

Maternal child health professionals can play a major role in a woman's infant feeding decision and in her success. Women in low-income ethnically diverse populations are especially at risk of not being counseled to breastfeed and not receiving sufficient follow-up.<sup>187</sup> In general, studies in the 1980s and 90s demonstrated that nurses breastfeeding knowledge was low<sup>25,188, 189</sup>, and health professionals were perceived as providing the least support in the form of encouragement, information, and assistance with breastfeeding techniques.<sup>190,191</sup> Issues of modeling bottle feeding in the hospitals, including the use of formula discharge packs, affected women's perceptions of health professionals' support of breastfeeding.<sup>192, 193</sup>

More recent studies indicate that many health professionals serving mothers and infants are inadequately prepared to provide prenatal education, perinatal support and postpartum follow-up for breastfeeding women. Doctors, nurses and other health care providers typically receive little education about lactation during their training.<sup>194-197</sup> A survey of 1137 pediatric fellows indicated the physicians were significantly lacking knowledge in the area of breastfeeding management.<sup>198</sup> An objective rating of lactation

content in 7 pediatric textbooks demonstrated that when breastfeeding information was not omitted, it was often highly variable, inconsistent and inaccurate.<sup>199</sup>

Specific to obstetric and pediatric visits, mothers reported minimal discussion of breastfeeding with their practitioners, and few reported discussing specific techniques to maintain lactation while returning to work.<sup>71</sup> Similarly, practitioners (medical doctors, nurse practitioners and nurse midwives) identified limited time during visits as a barrier to breastfeeding promotion.<sup>200</sup> Obstetric providers were least confident about managing insufficient milk supply while pediatric providers were not confident in dealing with breast pain and cracked nipples. Pediatric providers tend to recommend supplementation when infants are slow to gain weight.<sup>200</sup> The unintentional gaps in information, the neutral attitude, and lack of collaboration with breastfeeding specialists can have a significant effect on the quality of breastfeeding promotion and support provided to women and children.<sup>71, 200-201</sup>

Research supports the need for health care professionals to continue to examine the breastfeeding-related messages they project to mothers, verbal as well as nonverbal. Promotion and support efforts should include mothers' significant others, including fathers and maternal grandmothers.<sup>202-205</sup> Health care professionals need further education on lactation management, especially in the areas of cultural practices, medication use in lactating women, and managing breastfeeding and employment.<sup>207-210</sup> Training of health professionals has been shown in some studies to have an effect on practice and breastfeeding rates.<sup>211-214</sup> Further research is needed to identify effective methods of education to empower health care providers in their efforts to support their breastfeeding patients through collaboration, leadership and practice.

## **Breastfeeding practices in developing countries: insights from Africa**

Breastfeeding is almost universally practiced in most developing countries. In Africa approximately 99% of mothers breastfeed their infants (Figure 1). Breast milk is typically regarded as the best food for an infant. According to the DHS (Demographic and Health Surveys) comparative studies series 30,<sup>215</sup> there were no significant differences in the proportion of children ever breastfed when breastfeeding rates were examined by socio-and bio demographic variables. These variables include mothers' education, rural/urban residence, mother's age, birth order, sex of child, birth multiplicity and birth interval. In a study in Botswana in 2002, 95% of the mothers interviewed had breastfed their infants.<sup>216</sup>

### Breast Feeding Patterns, Age of Introduction of Complementary Feeding

Although breast-feeding is almost universally practiced among mothers in most parts of the developing world, the predominant pattern is mixed feeding where an infant is given breast milk and other feeds such as water, infant formula, local and commercially prepared cereals, usually from birth. It is believed that newborns require water after food, in the same way that an adult also requires water after a meal. Most mothers supplement their milk with water, teas, or herbal preparations frequently from the first week of life. They do so believing that such fluids will relieve pain (especially colic and earache), prevent and treat common colds and constipation; soothe fretfulness and quench thirst.<sup>217</sup> Newborns are often given water from the family's normal source to introduce the child to the family's drinking water. Mothers are expected to rest after the "strenuous" process of labor, so glucose water is given to the baby to prevent hunger until lactation begins. It is believed that herbs will make the child "strong," cleanse the baby's stomach, or in other

cases, improve the child's health if the child appears to be ill. It has been shown however that many of these prelacteal feeds are harmful and increase the prevalence of diarrhea and respiratory infection.<sup>218,219</sup> In rural communities in the savannah area of Nigeria, breastfeeding is widely practiced, but prelacteal feeds are given by all mothers.<sup>220</sup>

### Breastfeeding Initiation

The association between the time of the first breastfeeding to subsequent breastfeeding establishment has been documented.<sup>201,202</sup> Earlier initiation of breastfeeding is associated with better breastfeeding outcomes. The range of percentage of infants' breastfed immediately after birth is between 10 (in Mali) to 44 (in Cote d'Ivoire) in West Africa.<sup>215</sup> Many countries have very low rates of immediate initiation of breastfeeding, though the majority is breastfed within the first 24 hours. Urban-Rural residence, maternal education and age only slightly affect of breastfeeding initiation (Table 2). Among survey respondents in rural southwestern Nigeria, 20% of mothers stated that breastfeeding should commence within 30 minutes of delivery, while about 10% said one or more days after delivery.<sup>227</sup> Three-quarters (76%) of the respondents would not give their newborn colostrum. Colostrum is believed to be bad milk that has stayed in the breast during pregnancy and as such, is harmful to the child and should be discarded. Apart from denying newborns the important benefits of colostrum, discarding it also delays initiation of breast-feeding.

The belief that breast milk alone is not sufficient to meet an infant's nutritional needs also leads to early supplementation. This is in line with child feeding practices in many parts of the world where early supplementation with water and other fluids is a common practice.<sup>207-208</sup>

### Demand feeding, expression of breast milk and “wet-nursing”

Breastfeeding on demand is believed to be the normal way of feeding infants. In a survey in Western Nigeria, <sup>227</sup>75% of mothers believed in demand feeding (i.e. a baby should be offered breast milk when he or she needs it, when the baby cries or appears to be hungry). Mothers were aware that some mothers “in the city” gave their infants breast milk based on time intervals, but they thought this was unnatural; and wrong. Mothers in rural areas are more likely to breastfeed on demand. Similar observations have also been made among the Bambaras of Mali. <sup>209</sup>

The use of expressed breast milk is not culturally accepted in many parts of Africa. If a mother has to go out, for example to work, it is believed that her child should be fed supplements till she returns. Body fluids (blood, breast milk) and other parts (hair, nails) can be used to divine illness against their owners. Therefore, expressed breast milk can be used to poison, harm, or bewitch a mother or her infant. The milk can also go sour. <sup>227</sup> When an infant cannot be breastfed by its mother, either as a result of death or illness, the use of a wet-nurse is recommended. This wet-nurse is usually an older woman in the family. She can be the sister of the child’s father, or a co-wife of the mother. The most important thing is that the mother of the child and the wet nurse must have had an amicable relationship, and their “blood can mix well.”<sup>210</sup>

### Duration of breastfeeding

The age at which breastfeeding is terminated varies between one to two years. In sub-Saharan Africa, the median duration of breastfeeding, overall, is about 21 months. <sup>199</sup> Rural children are breast fed 3.2 months longer than urban children. (Table 3) Rural mothers are usually housewives, farmers, or self-employed, and thus have their children

with them most of the time; moreover infant formula may not be readily accessible or affordable. Mothers in the urban areas on the other hand, are more likely to work outside the home leaving their children in day care centers or with housemaids. They may also have more access and money to purchase infant formula and other food supplements.

Breastfeeding is often terminated if the lactating mother becomes pregnant. Pregnant women are not expected to breastfeed as it is believed that the child and the fetus could be harmed. To prevent this, breastfeeding is terminated and the child is washed with an herbal preparation.<sup>210</sup>

## Conclusion

In the last 20 years the knowledge base from which lactation consultants can draw from has grown phenomenally. Methods of research have also grown in caliber and sophistication. Knowledge of the physiologic processes of lactogenesis and the maintenance of lactation has progressed significantly, as has the epidemiologic evidence of the benefits of breastfeeding and human milk for the infant and mother. In addition, we have learned about the social, cultural, and personal context surrounding and impacting the complex process of initiation and continuation of breastfeeding. The impact of health care professionals on promoting and supporting breastfeeding has improved, but more is needed in our medical and nursing educational preparation to improve care of the breastfeeding mothers. Methods of treating common problems in breastfeeding mothers have progressed but more research is needed relative to prevention of serious problems such as insufficient milk syndrome and non-exclusive breastfeeding. Breastfeeding among vulnerable populations has increased in general, but more rigorous clinical trials

are needed to promote breastfeeding in women of color, teen mothers, and mothers of very low birth weight infants. As our knowledge of the biological, behavioral and environmental factors that affect breastfeeding continues to grow, lactation consultants will identify additional research problems and areas of needed knowledge. Thus the cycle will continue of describing and explaining problems and ultimately testing interventions to improve practice. It is through this cycle that promotion and support of breastfeeding will continue and likewise improve breastfeeding initiation and duration worldwide.

Table 1.

Function	Key Constituents
Anti-microbial	Oligosaccharides, lactoferrin, fatty acids, lysozyme, immunoglobulins, bifidus factor, complement, mucins, lactoperoxidase
Anti-inflammatory	Cytokines, Long-chain polyunsaturated fatty acids, growth factors, lactoferrin, hormones
Immune system promotion/development	Macrophages, cytokines, lymphocytes, long-chain polyunsaturated fatty acids, nucleotides, growth factors, hormones, neutrophils

**Table 2: Regional Summary of Timing of Initial Breastfeeding**

Region	Timing of initial breastfeeding after birth				
	First hour	1-23 hours	2 <sup>nd</sup> day	After 2days	Total
Sub-Saharan Africa	32.5	31.9	14.3	21.3	100.0
Near East/North Africa	35.6	40.5	9.7	14.2	100.0
Asia	15.0	31.6	16.5	36.9	100.0
Latin America/Caribbean	36.6	29.1	13.2	21.2	100.0

Source: demographic and Health surveys, 1990-96<sup>1</sup>



**Table 3 Regional summary of median duration of breastfeeding (in months), by urban-rural residence**

Region	Residence		
	Total	Urban	Rural
Sub-Saharan Africa	21.0	18.4	21.6
Near East/ North Africa	14.8	13.3	15.3
Asia	21.3	18.9	22.0
Latin America/Caribbean	13.9	12.7	15.0

Source: demographic and Health surveys, 1990-96<sup>1</sup>

#### References

1. Sackett DL, Rosenberg WMC, Gray JAM, Haynes RB, Richardson WS. Evidence based medicine: what it is and what it isn't. *BMJ*. 1996;312: 71-72.
2. Cook TD, Campbell DT. *Quasi-experimentation: Design and analysis issues*. Boston: Houghton-Mifflin; 1979.
3. Fawcett J. *The Relationship of Theory And Research (3<sup>rd</sup> edition)*. Philadelphia: Davis;1999.
4. Labbok MH, Coffin CJ. A call for consistency in definition of breastfeeding behaviors. *Social Science & Medicine*. 1997; 44: 1931-2.
5. Heinig MJ. Host defense benefits of breastfeeding for the infant. *Pediatr Clin North Am*. 2001; 48: 105-123.
6. Field CJ. The immunological components of human milk and their effect on immune development in infants. *J Nutr* 2005; 135: 1-4.

7. Kelleher SL, Lonnerdal B. Immunological activities of associated with milk. *Adv Nutr Res.* 2001; 10: 39-65.
8. Hartmann PE, Cregan MD, Mitoulas LR. Maternal modulation of specific and non-specific immune compounds of colostrum and mature milk. *Adv Nutr Res.* 2001; 365-87.
9. Perez-Escamilla, R., & Guerrero, M. L. Epidemiology of breastfeeding: advances and multidisciplinary applications. *Advances in Experimental Medicine & Biology*, 2004;554: 45-59.
10. Heinig MJ, Dewey KG. Health advantages of breastfeeding for infants: a critical review. *Nutr Res Rev.* 1996; 9:89-110.
11. Kramer MS, Kakuma R. The optimal duration of exclusive breastfeeding: a systematic review. *Adv Exp Med Biol.* 2004; 554: 63-77.
12. Gartner LM, Morton J, Lawrence RA et al. American Academy of Pediatrics. Breastfeeding and the use of human milk. *Pediatrics* 2005; 115: 496-506.
13. Labbok, M., & Krasovec, K. Toward consistency in breastfeeding definitions. *Studies in Family Planning*, 1990;21: 226-230.
14. World Health Organization. Global Data Bank on Breastfeeding. Geneva, Switzerland: World Health Organization; 1996
15. Callen J, Pinelli J. A review of the literature examining the benefits and challenges, incidence and duration, and barriers to breastfeeding in preterm infants. *Adv Neonatal Care.* 2005; 5: 72-88.
16. Anderson JW, Johnstone BM, Remley DT. Breastfeeding and cognitive development: a meta-analysis. *Am J Clin Nutr* 1999; 70: 525-535.
17. Feldman R, Eidelman AI. Direct and indirect effects of breast milk on the neurobehavioral and cognitive development of premature infants. *Dev Psychobiol.* 2003; 43: 109-119.
18. McGuire W, Anthony MY. Donor human milk versus formula for preventing necrotizing enterocolitis in preterm infants: systemic review. *Arch Dis Child Fetal Neonatal Ed.* 2003; 88: F11-14.
19. De Silva A, Jones PW, Spencer SA. Does human milk reduce infection rates in preterm infants? A systemic review. *Arch Dis Child Fetal Neonatal Ed.* 2004; 89: F509-13.

20. Davis MK. Breastfeeding and chronic disease in childhood and adolescence. *Pediatrics Clinics of North America* 2001; 48: 125-142.
21. Muraro A, Dreborg S, Halken, S et al. Dietary prevention of allergic diseases in infants and small children. Part III. Critical review of published peer-reviewed observational and interventional studies and final recommendations. *Pediatr Allergy Immunol.* 2004; 15: 291-307.
22. Oddy WH, Peat JK. Breastfeeding asthma, and atopic disease: an epidemiological review of the literature. *J Hum Lact.* 2003; 19: 250-261.
23. Dewey KG. Does breastfeeding protect against childhood obesity? *J Hum Lact.* 2003; 19: 9-18.
24. Horwood LJ, Fergusson DM. Breastfeeding and later cognitive and academic outcomes. *Pediatrics* 1998; 101: E9.
25. Anderson E, Geden E. Nurses' knowledge of breastfeeding. *J Obstet, Gynecol, and Neonatal Nurs* 1991;20:58-64.
26. Jain A, Concato J, Leventhal JM. How good is the evidence linking breastfeeding and intelligence? *Pediatrics* 2002; 109: 1044-1053.
27. Labbok MH. Effects of breastfeeding on the mother. *Pediatrics Clinics of North America* 2001; 48: 143-158.
28. Heinig MJ, Dewey KG. Health effects of breastfeeding for mothers: a critical review. *Nutr Res Rev.* 1997; 10: 35-56.
29. Collaborative Group on Hormonal Factors in Breast Cancer. Breast cancer and breastfeeding: collaborative reanalysis of individual data from 47 epidemiological studies in 30 countries, including 50302 women with breast cancer and 96973 women without the disease. *Lancet* 2002; 360: 187-195.
30. Riman T, Nilsson S, Persson IR. Review of epidemiological evidence for reproductive and hormonal factors in relation to the risk of epithelial ovarian malignancies. *Acta Obstet Gynecol Scand.* 2004; 83: 783-795.
31. Callen, J., & Pinelli, J. Incidence and duration of breastfeeding for term infants in Canada, United States, Europe and Australia: A literature review. *Birth.* 2004; 31: 285-292.
32. Duckett, L., Henly, S., Avery, M., Potter, S., Hills-Bonczyk, S., Hulden, R., et al. A theory of planned behavior-based structural model for breast-feeding. *Nursing Research,* 1998; 47: 325-335.

33. Guise, J. M., Palda, V., Westhoff, C., Chan, B. K. S., Helfand, M., & Lieu, T. A. The effectiveness of primary care-based interventions to promote breastfeeding: systematic evidence review and meta-analysis for the US Preventive Services Task Force. *Annals of Family Medicine*, 2003; 1: 70-78.
34. Kronborg, H., & Vaeth, M. The influence of psychosocial factors on the duration of breastfeeding. *Scandinavian Journal of Public Health*, 2004; 32: 210-206.
35. McInnes RJ, Stone DH. The process of implementing a community-based peer breastfeeding support programme: the Glasgow experience. *Midwifery*. 2001; 17: 65-73.
36. Nagy E, Orvos H, Pal A, Kovacs L, Loveland K. Breastfeeding duration and previous breastfeeding experience. *Acta Paediatr*. 2001; 90: 51-60.
37. Hill, P. D. Update on breastfeeding: Healthy People Objectives. *MCN*. 2000; 25: 248-251.
38. Stewart-Knox B, Gardiner K, Wright M. What is the problem with breastfeeding? A qualitative analysis of infant feeding perceptions. *J Hum Nutr Diet*. 2003; 16: 217-218.
39. Rea MF, Morrow AL. Protecting, promoting, and supporting breastfeeding among women in the labor force. *Adv Exp Med Biol*. 2004; 554: 121-132.
40. Butler S, Williams M, Tukuitonga C, Paterson J. Factors associated with not breastfeeding exclusively among mothers of a cohort of Pacific infants in New Zealand. *N Z Med J*. 2004; 117: U908.
41. Cattaneo A, Yngve A, Loletzko B, Guzman LR. Protection, promotion and support of breastfeeding in Europe: current situation. *Public Health Nutrition*. 2005; 8: 39-46.
42. Dodgson, J. E., Henly, H. J., Duckett, L., & Tarrant, M. Theory of planned behavior-based models for breastfeeding duration among Hong Kong mothers. *Nursing Research*, 2002; 52: 148-158.
43. Perez-Escamilla, R., Cobas, J. A., Balcazar, H., & Benin, M. H. Specifying the antecedents of breast-feeding duration in Peru through a structural equation model. *Public Health Nutrition*, 1999; 2: 461-467.
44. Rogers, I. S., Emmett, P. M., & Golding, J. The incidence and duration of breast feeding. *Early Human Development*, 1997; 49: S45-74.
45. Dennis CL. Breastfeeding Initiation and Duration: A 1990-2000 Literature Review. *J Obstet, Gynecol, and Neonatal Nurs* 2002;31:12-32.
46. Li, R., Fridinger, F., & Grummer-Strawn, L. Public perceptions on breastfeeding constraints. *Journal of Human Lactation*, 2002; 18: 227-235.

47. Scott JA, Landers MC, Hughes RM, Binns CW. Psychosocial factors associated with the abandonment of breastfeeding prior to hospital discharge. *J Hum Lact*. 2001; 17: 24-30.
48. McIntyre E, Hiller JE, Turnbull D. Community attitudes to infant feeding. *Breastfeed Rev*. 2001; 9: 27-33.
49. Jelliffe, D. B., & Jelliffe, P. E. F. *Human Milk in the Modern World: Psychosocial, Nutritional and Economic Significance*. Oxford: Oxford University Press. 1978.
50. Shaw, E., & Kaczorowski, K. The effect of a peer counseling program on breastfeeding initiation and longevity in a low-income rural population. *J Hum Lact*, 1999; 15:19-25.
51. Zimmerman, D. R. You can make a difference: Increasing breastfeeding rates in an inner-city clinic. *J Hum Lact*, 1991; 15:217-220.
52. Kang NM, Song Y, Hyun TH, Kim KN. Evaluation of the breastfeeding intervention program in a Korean community health center. *Int J Nurs Stud*. 2005; 42: 409-413.
53. Chapman DJ, Damio G, Young S, Perez-Escamilla R. Effectiveness of breastfeeding peer counseling in a low-income, predominantly Latina population. *Obstet Gynecol Surv*. 2005; 60: 157-158.
54. Forster D, McLachlan H, Lumley J, Beanland C, Walderstrom U, Amir L. Two mid-pregnancy interventions to increase the initiation and duration of breastfeeding: a randomized controlled trial. *Birth*. 2004; 31: 176-82.
55. Quinlivan JA, Box H, Evans SF. Postnatal home visits in teenage mothers: a randomized controlled trial. *Lancet*. 2003; 361: 893-900.
56. Lu, M. C., Prentice, J., Yu, S. M., Inkelas, M., Lange, L. O., & Halfon, N. Childbirth education classes: Sociodemographic disparities in attendance and the association of attendance with breastfeeding initiation. *Maternal and Child Health Journal*, 2003; 7: 87-93.
57. Chatterji, P., & Brookes-Gunn, J. WIC participation, breastfeeding practices and well-child care among unmarried, low-income mothers. *American Journal of Public Health*, 2004; 94, 1324-1327.
58. Ahluwalia, I. B., Morrow, B., Hsia, H., & Grummer-Strawn, L. M. Who is breastfeeding? Recent trends from the Pregnancy risk assessment and monitoring system. *Journal of Pediatrics*. 2003;142:486-491.
59. WHO Global Databank. World Health Organization. URL: [http://www.who.int/nut/db\\_bfd.htm](http://www.who.int/nut/db_bfd.htm), accessed 5/24/05.

60. Bland, R. M., Rollins, N. C., Solarsh, G., Broeck, J. V. d., & Coovadia, H. M. Maternal recall of exclusive breast feeding duration. *Archives of Disease in Childhood*. 2003; 88: 778-783.
61. Li, R., Ballew, C., Gillespie, K., Grummer-Strawn, L., & Ogden, C. Prevalence of exclusive breastfeeding among US infants: The third National Health and Nutrition Examination Survey (Phase II, 1991-1994). *American Journal of Public Health*, 2002; 92: 1107-1011.
62. Dubois, L., & Girard, M. Social determinants of initiation, duration and exclusivity of breastfeeding at the population level: The results of the Longitudinal Study of Child Development in Quebec (ELDEQ 1998--2002). *Canadian Journal of Public Health*, 2003; 94: 300-305.
63. Bernardo, H., Kramer, M. S., & Platt, R. W. Maternal smoking and the risk of early weaning: A meta-analysis. *American Journal of Public Health*. 2001; 91: 304-307.
64. Blyth R, Creedy DK, Dennis C, Moyle W, Pratt J, De Vries SM. Effect of maternal confidence on breastfeeding duration: an application of breastfeeding self-efficacy theory. *Birth*. 2002 ; 29: 278-4
65. Janke, J. R. Prediction of breast-feeding attrition: Instrument development. *Applied Nursing Research*, 1992; 5: 48-53.
- 66 Dogson JE, Henly HJ, Duckett L, Tarrant M. Theory of planned behavior-based models for breastfeeding duration among Hong Kong mothers. *Nurs Res*. 2003; 52: 148-158.
67. Sikorski, J., Renfrew, M. J., Pindoria, S., & Wade, A. (2005). Support for breastfeeding mothers. *The Cochrane Database of Systematic Reviews*, 1, 1-14.
68. Martens, P. J. Increasing breastfeeding initiation and duration at a community level: An evaluation of Sagkeeng First Nation's community health nurse and peer counselor programs. *Journal of Human Lactation*, 2002; 18: 236-246.
69. Pugh, L. C., Milligan, R. A., Frick, K. D., Spatz, D., & Bronner, Y. Breastfeeding duration, cost and benefits of a support program for low-income breastfeeding women. *Birth*, 2002; 29: 95-100.
70. Dearden, K., Altaye, M., Maza, I. d., Oliva, M. d., Stone-Jimenez, M., Burkhalter, B. R., et al. The impact of mother-to-mother support on optimal breast-feeding: A controlled community intervention trial in peri-urban Guatemala City, Guatemala. *Pan American Journal of Public Health*, 2002; 12: 193-201.

71. Taveras EM, Li, R., Grummer-Strawn, L., Richardson, M., Marshall, R., Rego, V.H., Miroshnik, I., & Lieu, T.A. Mothers' and clinicians' perspectives on breastfeeding counseling during routine preventive visits. *Pediatrics* 2004;113:e-405-11.
72. Neville MC. Anatomy and physiology of lactation. *Pediatrics Clinics of North America* 2001; 48: 13-34.
73. Russo J, Russo IH. Development of the human breast. *Maturitas*. 2004;49:2-15.
74. Buhimschi CS. Endocrinology of lactation. *Obstet Gynecol Clin North Am*. 2004; 31: 963-979.
75. Daly SE, Hartmann PE. Infant demand and milk supply. Part 1. Infant demand and milk production in lactating women. *J Hum Lact* 1995; 11: 21-26.
76. Daly SE, Hartmann PE. Infant demand and milk supply. Part 2. The short-term control of milk synthesis in lactating women. *J Hum Lact*. 1995; 11: 27-27.
77. Rigg LA, Yen SSC. Multiphasic prolactin secretion during parturition in human subjects. *Am J obstet Gynecol* 1977; 128: 215.
78. Sala-Vila A, Castellote AI, Rodriguez-Palmero M, Campoy C, Lopez-Sabater MC. Lipid composition in human breast milk from Granada (Spain): Changes during lactation. *Nutrition*. 2005;21:467-73.
79. Neville MC, Morton J, Unemura S. Lactogenesis: Transition from pregnancy to lactation. *Pediatrics Clinics of North America* 2001; 48: 35-52.
80. Dewey KG. Maternal and fetal stress are associated with impaired lactogenesis in humans. *J Nutr* 2001; 131: 3012S-3015S.
81. Dewey KG, Nommsen-Rivers LA, Heinig MJ, Cohen RJ. Risk factors for suboptimal infant breastfeeding behavior, delayed onset of lactation, and excess neonatal weight loss. *Pediatrics*. 2003; 112: 607-619.
82. Bailey VF, Sherriff J. Reasons for the early cessation of breastfeeding in women from lower socio-economic groups in Perth, Western Australia. *Breastfeeding Rev*. 1993; 27: 390-3.
83. Bourgoin GL, Lahaie NR, Rheaume BA, Berger MG, Dovigi CV, Picard LM, Sahai VF. Factors influencing the duration of breastfeeding in the Sudbury Region. *Can J Pub Health*. 1997; 88: 238-41.
84. Hill PD, Humenick SS. Nipple pain during breastfeeding: the first two weeks and beyond. *J Perinatal Education*. 1993; 2: 21-35

85. Ziemer MM, Pigeon JG. Skin changes and pain in the nipple during the 1st week of lactation. *J Obstet Gynecol Neonatal Nurse*. 1993; 22: 247-56.
86. Blair A, Cadwell K, Turner-Maffei C, Brimdyr K. The relationship between positioning, the breastfeeding dynamic, the latching process and pain in breastfeeding mothers with sore nipples. *Breastfeeding Rev*. 2003;11:5-10.
87. Woolridge MW. Aetiology of sore nipples. *Midwifery*. 1986; 2: 172-6.
88. Righard L. Are breastfeeding problems related to incorrect breastfeeding technique and the use of pacifiers and bottles? *Birth*. 1998; 25: 40-4.
89. Amir LH, Harris H, Andriske L. An audit of mastitis in the emergency department. *J Hum Lact*. 1999;15:221-224.
90. Marmet C, Shell E, Marmet R. Neonatal frenotomy may be necessary to correct breastfeeding problems. *J Hum Lact*. 1990; 6: 117-21.
91. Wilton JM. Sore nipples and slow weight gain related to a short frenulum. *J Hum Lact*. 1990; 6: 122-3.
92. Centuori S, Burmaz T, Ronfani L, Fragiaco M, Quintero S, Pavan C, Davanzo R, Cattaneo A. Nipple care, sore nipples, and breastfeeding: a randomized trial. *J Hum Lact*. 1999; 15: 125-30.
93. Pugh LC, BL, Buchko, Bishop BA, Cochran JF, Smith LR, Lerew DJ. A comparison of topical agents to relieve nipple pain and enhance breastfeeding. *Birth*. 1996; 23: 88-93.
94. Dodd, V. and C. Chalmers. Comparing the use of hydrogel dressings to lanolin ointment with lactating mothers. *J Obstet Gynecol Neonatal Nurse*. 2003; 32; 486-94.
95. Brent N, Rudy SJ, Redd B, Rudy TE, Roth LA. Sore nipples in breast-feeding women: a clinical trial of wound dressings vs conventional care. *Arch Pediatrics & Adolescent Medicine*. 1998; 152: 1077-82.
96. Lavergne NA. Does application of tea bags to sore nipples while breastfeeding provide effective relief? *J Obstet, Gynecol Neonatal Nurse*. 1997; 26: 53-8.
97. Humenick SS, Hill PD, Anderson MA. Breast engorgement: patterns and selected outcomes *J Hum Lact*. 10: 87-93.
98. Moon JL, Humenick SS. Breast engorgement: contributing variables and variables amenable to nursing intervention. *J Obstet Gynecol Neonatal Nurse*. 1989;18: 309-15.
99. Snowden HM, Renfrew MJ, Woolridge MW. Treatments for breast engorgement during lactation. *The Cochrane Library (Oxford)*. 2005;(1)



100. Fetherston C. Characteristics of lactation mastitis in a Western Australian cohort. *Breastfeeding Rev.* 1997;5:5-11.
101. Foxman B, D'Arcy H, Gillespie B, Bobo JK, Schwartz K. Lactation mastitis: occurrence and medical management among 946 breastfeeding women in the United States. *Am J Epidemiol.* 2002;155:103-114.
102. Kinlay JR, O'Connell DL, Kinlay S. Incidence of mastitis in breastfeeding women during the six months after delivery: a prospective cohort study. *Med J Aust.* 1998;169:310-312.
103. Vogel A, Hutchison BL, Mitchell EA. Mastitis in the first year postpartum. *Birth.* 1999;26:218-225.
104. Fetherston C. Risk factors for lactation mastitis. *J Hum Lact.* 1998;14:109.
105. Osterman KL, Rahm V. Lactation mastitis: bacterial cultivation of breast milk, symptoms, treatment, and outcome. *J Hum Lact.* 2000;16:297-302.
106. Hager WD, Barton JR. Treatment of sporadic acute puerperal mastitis. *Infec Dis Obstet Gynecol.* 1996;4:97-101.
107. Matheson I, Aursnes I, Horgen M, Aobo O, Melby K. Bacteriological findings and clinical symptoms in relation to clinical outcome in puerperal mastitis. *Acta Obstet Gynecol Scand.* 1988; 67:723-726.
108. Fetherston C. Management of lactation mastitis in a Western Australian cohort. *Breastfeeding Rev.* 1997;5:13-19.
109. Wambach, K. A. Lactation mastitis: a descriptive study of the experience. *J Hum Lact.* 2003; 19: 24-34.
110. Binns CW, Scott JA. Breastfeeding: reasons for starting, reasons for stopping and problems along the way. *Breastfeeding Rev.* 2002; 10: 13-9.
111. Liamputtong P. Infant feeding practices: the case of Hmong women in Australia. *Health Care Women International.* 2002; 23: 33-48
112. McLennan JD. Early termination of breast-feeding in periurban Santo Domingo, Dominican Republic: mothers' community perceptions and personal practices. *Pan American J Public Health.* 2001; 9: 362-6.
113. Novotny R, Hla MM, Kieffer EC, Park C, Mor J, Thiele M. Breastfeeding duration in a multiethnic population in Hawaii. *Birth.* 2000; 27: 91-6.

114. Hill PD, Aldag J. Potential indicators of insufficient milk supply syndrome. *Res Nurs Health*. 1991; 14: 11-9.
115. Hill PD, Aldag JC Insufficient milk supply among Black and White breast-feeding mothers. *Res Nurs Health*. 1993; 16: 203-11.
116. Hill PD, Humenick SS. Development of the H & H Lactation Scale. *Nurs Research*. 1996; 45: 136-40.
117. Henly SJ, Anderson CM, Avery MD, Hills-Bonczyk SG, Potter S, Duckett LJ. Anemia and insufficient milk in first-time mothers. *Birth*. 1995; 22: 86-92.
118. Humenick SS, Hill PD, Thompson J, Hart AM. Breast-milk sodium as a predictor of breastfeeding patterns. *Can J Nurs Res*. 1998; 30: 67-81.
119. Neifert, M., S. DeMarzo, Seacat J, Young D, Leff M, Orleans M. The influence of breast surgery, breast appearance, and pregnancy-induced breast changes on lactation sufficiency as measured by infant weight gain. *Birth* 1990; 17: 31-8.
120. Willis CE, Livingstone V. Infant insufficient milk syndrome associated with maternal postpartum hemorrhage. *J Hum Lact*. 1995;11:123-6.
121. Yamauchi, Y. and I. Yamanouchi. The relationship between rooming-in/not rooming-in and breastfeeding variables. *Breastfeeding Rev*. 1992; 2: 238-41.
122. McCarter-Spaulding DE, Kearney MH. Parenting self-efficacy and perception of insufficient breast milk. *J Obstet Gynecol Neonatal Nurse*. 2001; 30: 515-22.
123. ILCA, First 14 days.
124. Hurst NM, Meier, PP 2005. Breastfeeding the preterm infant. *Breastfeeding and Human Lactation (3<sup>rd</sup> edition)*. Riordan, J. pp367-406
125. Hurst NM, Meier PP, Engstrom JL, Myatt A. Mothers performing in-home measurement of milk intake during breastfeeding of their preterm infants: maternal reactions and feeding outcomes. *J Hum Lact*. 2004; 20:178-87.
126. Kavanaugh K, Mead L, Meier P, Mangurten HH Getting enough: mothers' concerns about breastfeeding a preterm infant after discharge *J Obstet Gynecol Neonatal Nurse* 1995; 24: 23-32.
127. Kavanaugh K, Meier P, Zimmermann B, Mead L. The rewards outweigh the efforts: breastfeeding outcomes for mothers of preterm infants. *J of Hum Lact* 1997; 13: 15-21.

128. Miracle DJ, Meier PP, Bennett PA. Mothers' decisions to change from formula to mothers' milk for very-low-birth-weight infants. *J Obstet Gynecol Neonatal Nurse*. 2004; 33: 692-703.
129. Meier P. Bottle- and breast-feeding: effects on transcutaneous oxygen pressure and temperature in preterm infants. *Nurs Research*. 1988; 37:36-41.
130. Meier P, Anderson GC. Responses of small preterm infants to bottle- and breast-feeding. *Am J Maternal Child Nurse* 1987;12: 97-105.
131. Meier P, Pugh EJ. Breast-feeding behavior of small preterm infants. *Am J Maternal Child Nurse*. 1985; 10:396-401.
132. Meier P, Wilks S. The bacteria in expressed mothers' milk. *Am J Maternal Child Nurse*. 1987; 12:420-3.
133. Hill PD, JC Aldag, Chatterton RT. Effects of pumping style on milk production in mothers of non-nursing preterm infants. *J Hum Lact*. 1999; 15: 209-16.
134. Hill PD, JC Aldag, Chatterton RT (2001). Initiation and frequency of pumping and milk production in mothers of non-nursing preterm infants. *J Hum Lact* 2001; **17**: 9-13.
135. Valentine CJ, Hurst NM, Schanler RJ. Hindmilk improves weight gain in low-birth-weight infants fed human milk. *J Pediatr Gastroenterol Nutr*. 1994; 18:474-7.
136. Meier PP, Brown LP, Hurst NM, Spatz DL, Engstrom JL, Borucki LC, Krouse AM. Nipple shields for preterm infants: effect on milk transfer and duration of breastfeeding. *J Hum Lact*. 2000; 16:106-14; 129-31.
137. Hill PD, Aldag JC, Chatterton RT, Zinaman M. Comparison of milk output between mothers of preterm and term infants: the first 6 weeks after birth. *J Hum Lact*. 2005; 21: 22-30.
138. Charpak N, Ruiz-Pelaez JG, Figueroa de C Z, Charpak Y. Kangaroo mother versus traditional care for newborn infants less than or equal to 2000 grams: a randomized, controlled trial. *Pediatrics*. 1997; 100: 682-8.
139. Hurst NM, Valentine CJ, Renfro L, Burns P, Ferlic L. Skin-to-skin holding in the neonatal intensive care unit influences maternal milk volume. *J Perinatology*. 1997; 17:213-7.
140. Ludington-Hoe SM, Thompson C, Swinth J, Hadeed AJ, Anderson GC. Kangaroo care: research results, and practice implications and guidelines... findings of two research projects. *Neonatal Network: The J of Neonatal Nursing*. 1994; 13: 19-27, 29-34.

141. Nyqvist KH, Rubertsson C, Ewald U, Sjoden P. Development of the Preterm Infant Breastfeeding Behavior Scale (PIBBS): a study of nurse-mother agreement. *J Hum Lact.* 1996; 12: 207-19.
142. Hurst NM, Myatt A, Schanler RJ. Growth and development of a hospital-based lactation program and mother's own milk bank. *J Obstet Gynecol Neonatal Nurse.* 1998; 27: 503-10.
143. Meier PP, Engstrom JL, Mangurten HH, Estrada E, Zimmerman B, Kopparthi R. Breastfeeding support services in the neonatal intensive-care unit. *J Obstet Gynecol Neonatal Nurse.* 1993; 22:338-47.
144. Meier PP, Engstrom JL, Mingoelli SS, Miracle DJ, Kiesling S. The Rush Mothers' Milk Club: breastfeeding interventions for mothers with very-low-birth-weight infants. *J Obstet Gynecol Neonatal Nurse.* 2004; 33:164-74.
145. Spear HJ. Nurses' attitudes, knowledge, and beliefs related to the promotion of breastfeeding among women who bear children during adolescence. *J Pediatric Nurs: Nurs Care of Children and Families.* 2004; 19: 176-83.
146. Brasil AL, Vitolo MR, Lopez FA, Nobrega FJ. Fat and protein composition of mature milk in adolescents. *J Adol Health,* 1991;12: 365-371.
147. Lipsman S, Dewey KG, Lönnerdal, B. Breast-feeding among teenage mothers: Milk composition, infant growth, and maternal dietary intake. *J Pediatric Gastroente Nutrition,*1985;\_4: 426-434
148. Motil KJ, Kertz B, Thotathuchery M. Lactational performance of adolescent mothers shows preliminary differences from that of adult women. *J Adol Health.* 1997; 20: 442-449.
149. Vitolo MR, Brasil AL, Lopez FA, Nobrega, J. Colostrum composition in adolescent mothers. *J Am Coll Nutr,*1993;12: 547-550.
150. Strobino DM, Ensminger ME, Nanda J, Kim YJ. Young motherhood and infant hospitalization during the first year of life. *J Adol Health,* 1992; 13: 553-560.
151. Wilson MD, Duggan AK, Joffe A. Rehospitalization of infants born to adolescent mothers. *J Adol Health Care,* 1990; 11, 510-515.
152. Hannon PR, Willis SK, Bishop-Townsend V, Martinez I, Scrimshaw S. African American and Latina adolescent mothers' infant feeding decisions and practices: A qualitative study. *J Adolescent Health.* 2000;26: 399-407.
153. Harner HH, McCarter-Spaulding D. Teenage mothers and breastfeeding: Does paternal age make a difference? *J Hum Lact.* 2004; 20: 404-408.

154. Ineichen B, Pierce M, Lawrenson, R. (1997). Teenage mothers as breastfeeders: Attitudes and behaviors. *J Adolescence*, 1997; 20: 505-509.
155. Maehr, JC, Lizarraga JL, Wingard DL, Felice ME. A comparative study of adolescent and adult mothers who intend to breastfeed. *J Adolescent Health Care*. 1993; 1: 453-457.
156. Park YK, Meier ER, Song WO. Characteristics of teenage mothers and predictors of breastfeeding initiation in the Michigan WIC program in 1995. *J Hum Lact* 2003; 19: 50-6.
157. Robinson JB, Hunt AE, Pope J, Garner, B. Attitudes toward breastfeeding among adolescent mothers from a WIC population in northern Louisiana. *J Am Diet Assoc*, 1993; 93: 1311-1313.
158. Wiemann CM, DuBois JC, Berenson, A.B. Racial/ethnic differences in the decision to breastfeed among adolescent mothers. *Pediatrics*., 1998; 101: e11.
159. Wambach KA, Koehn M. Experiences of infant-feeding decision-making among urban economically disadvantaged pregnant adolescents. *J Adv Nurs*. 2004; 48: 361-370.
160. Benson S. Adolescent mothers' experience of parenting and breastfeeding - a descriptive study. *Breastfeeding Rev*. 1996; 4: 19-27.
161. Dykes F, Moran VH, Burt S, Edwards J. Adolescent mothers and breastfeeding: experiences and support needs -- an exploratory study. *J Hum Lact*. 2003; 19(4): 391-401
162. Swanson, N. Infant feeding patterns of teenage mothers in a small New England town. *International J SocFamily*. 1988; 18: 249-282.
163. Neifert M, Gray J, Gary N, Camp, B. Effect of two types of hospital feeding gift packs on duration of breast-feeding among adolescent mothers. *J Adolescent Health Care*. 1988; 9: 411-413.
164. Volpe EM, Bear M. Enhancing breastfeeding initiation in adolescent mothers through the Breastfeeding Educated and Supported Teen (BEST) Club. *J Hum Lact*. 2000; 16, 196-200.
165. Li, L., Zhang, M., Scott, J. A., & Binns, C. W. Factors associated with the initiation and duration of breastfeeding by Chinese mothers in Perth, Western Australia. *J Hum Lact*, 2004; 20: 188-195.
166. Earle, S. Factors affecting the initiation of breastfeeding: Implications for breastfeeding promotion. *Health Promotion International*, 2002; 17: 205-214.

167. Tarrant, M., & Dodgson, J. E. Initiating and sustaining breastfeeding in Hong Kong: contextual influences on new mothers' experiences. *Nursing and Health Sciences*, 2002; 4: 189-199.
168. Dodgson, J. E., Duckett, L., Garwick, A., & Graham, B. L. An ecological perspective of breastfeeding within an indigenous community. *Journal of Nursing Scholarship*, 2002; 34: 235-241.
169. Kannan, S., Carruth, B. R., & Skinner, J. Cultural influences on infant feeding beliefs of mothers. *J Am Diet Assoc*, 1999; 99: 88-90.
170. Wright, A. L., Naylor, A., Wester, R., Bauer, M., & Sutcliffe, E. Using cultural knowledge in health promotion: Breastfeeding among the Navajo. *Health Education & Behavior*, 1997; 24, 625-639.
171. Homer CS, Sheehan A, Cooke M. Initial infant feeding decisions and duration of breastfeeding in women from English, Arabic, and Chinese-speaking backgrounds in Australia. *Breastfeed Rev*. 2002; 10: 27-32.
172. Denman-Vitale S & Murillo EK. Effective promotion of breastfeeding among Latin American women newly immigrated to the United States. *Holist Nurs Pract*. 1999;3: 51-60.
- 173 Campbell H, Jones I. Promoting breastfeeding: a view of the current position and a proposed agenda for action in Scotland. *J Pub Health Med*. 1996; 18: 406-414.
174. Li R, Zhao A, Mokdad A, Barker L, Grummer-Strawn L. Prevalence of breastfeeding in the United States: The 2001 National Immunization Survey. *Pediatrics* 2003;111:1198-1201.
175. Guttman N, & Zimmerman, D.R. Low-income mothers' views on breastfeeding. *Soc Sci and Med* 2000;50:1457-1473.
176. Timbo B, Altekruze S, Headrick M, Klontz K. Breastfeeding among black mothers: evidence supporting the need for prenatal intervention. *J of Soc Pediatric Nurs* 1996;1:35-40.
177. Balcazar H, Trier CM, & Cobas JA. What predicts breastfeeding intention in Mexican-American and non-Hispanic white women? Evidence from a national survey. *Birth*. 1995; 22: 74-80.
178. Rassin DK, Markides KS, Baranowski T, Richardson CJ, Mikrut WD, & Bee D. 1993 Acculturation and initiation of breastfeeding. *J Clin Epidemiol*. 2000;47: 739-746.

179. Gill SL, Reifsnider E, Mann AR, Villarreal P, & Tinkle MB. Assessing Infant Breastfeeding Beliefs Among Low-Income Mexican Americans *Journal of Perinatal Education*. 2004;13: 1-12.
180. Sciacca JP, Dube DA, Phipps, BL & Ratliff MI. A breast feeding education and promotion program: Effects on knowledge, attitudes, and support for breast feeding. *Journal of Community Health*. 1995;20: 473-490.
181. Lazarov M & Evans A. Breastfeeding—Encouraging the best for low-income women. *Zero to Three*. 2000; 15-23.
182. McInnes RJ, Love JG, Stone DH. Independent predictors of breastfeeding intention in a disadvantaged population of pregnant women. *BMC Public Health*. 2001; 1: 10.
183. Avery M, Duckett L, Dodgson J, Savik K, Henly SJ. Factors associated with very early weaning among primiparas intending to breastfeed. *Maternal and Child Health Journal*. 1998; 2: 167-79
184. Bevan ML, Mosley, D., Lobach, K.S., Solimano, G.R. Factors influencing breast feeding in an urban WIC program. *J of the Amer Dietetic Assoc* 1984;84:563-567.
185. Dennis C. Breastfeeding peer support: Maternal and volunteer perceptions from a randomized controlled trial. *Birth*. 2002b;29: 169-176.
186. Scott JA, Landers MCG, Hughes RM, & Binns CW Factors associated with breastfeeding at discharge and duration of breastfeeding. *Journal of Paediatrics & Child Health*. 2001;37: 254-261.
187. Beal A, Kuhlthau K, Perrin J. Breastfeeding advice given to African American and white women by physicians and WIC counselors. *Public Health Reports* 2003; 118:368-376.
188. Crowder DS. Maternity nurses knowledge of factors promoting successful breastfeeding. *J Obstet, Gynecol, and Neonatal Nurs* 1981;10:28-30.
189. Hayes B. Inconsistencies among nurses in breast-feeding knowledge and counseling. *J Obstet, Gynecol, and Neonatal Nurs* 1981:430-433.
190. Albers RM. Emotional support for the breastfeeding mother. *Issues in Compr Pediatric Nurs* 1981;5:109-124.
191. Ellis DJ, Hewat RJ. Do we support breastfeeding mothers? *Midwives Chronicle and Nursing Notes* 1984;97:45-47.

192. Howard CR, Howard FM, Weitzman MC. Infant formula distribution and advertising in pregnancy: a hospital survey. *Birth: Issues in Perinatal Care and Education* 1994;21:14-19.
193. Reiff MI, Essock-Vitale SM. Hospital influences on early infant feeding practices. *Pediatrics* 1985;76:872-879.
194. Wong SC. Physicians should be educated on the benefits of breastfeeding. (Letters to the Editor). *Amer Fam Phys* 2002;66:209-210.
195. Freed G. National assessment of physician's breast-feeding knowledge, attitudes, training and experience. *JAMA* 1995;273:472-76.
196. Guise JMF, G. Resident physicians' knowledge of breastfeeding and infant growth. *Birth* 2000;27: 49-53.
197. Howard CR, Schaffer SJ, Lawrence RA. Attitudes, practices, and recommendations by obstetricians about infant feeding. *Birth* 1997;24:240-252.
198. Schanler RJ, O'Connor, K.G., Lawrence, R.A. Pediatrician's practices and attitudes regarding breastfeeding promotion. *Pediatrics* 1999;103:E35.
199. Philipp BL, Merewood A, Gerendas EJ, Bauchner H. Breastfeeding information in pediatric textbooks needs improvement. *J of Hum Lact* 2004;20(2):206-210.
200. Taveras EM, Li, R., Grummer-Strawn, L., Richardson, M., Marshall, R., Rego, V.H., Miroshnik, I., & Lieu, T.A. Opinions and practices of clinicians associated with continuation of exclusive breastfeeding. *Pediatrics* 2004;113:283-290.
201. DiGirolamo AM, Grummer-Strawn, L.M., & Fein, S.B. Do perceived attitudes of physicians and hospital staff affect breastfeeding decisions? *Birth* 2003;30(2):94-100.
202. Cornett B. Postpartal mothers' perceptions of in-hospital breastfeeding information and affective support as related to breastfeeding status.: OHIO STATE UNIVERSITY; 1989.
203. Beardshaw T. Supporting the role of fathers around the time of birth. *MIDIRS Midwifery Digest* 2001;11:476-479.
204. Barton S. Infant feeding practices of low-income rural mothers. *MCN: The Amer J of Mat/Child Nurs* 2001; 26:93-97.
205. Shaker I, Scott J, Reid R. Infant feeding attitudes of expectant parents: breastfeeding and formula feeding. *J of Adv Nurs* 2004;45:260-268.



206. Ekström A, Widström A, Nissen E. Breastfeeding support from partners and grandmothers: perceptions of Swedish women. *Birth* 2003;30:261-266.
207. Dykes F. Protecting, promoting and supporting breastfeeding. *British J of Midwifery* 2003;11(10):S24-8.
208. Shaw R, Wallace L, Bansal M. Is breast best? Perceptions of infant feeding. *Community Practitioner* 2003;76:299-303.
209. Jones W, Brown D. The medication vs breastfeeding dilemma. *British J of Midwifery* 2003;11(9):550-555.
210. Mlay R, Keddy B, Stern P. Demands out of context: Tanzanian women combining exclusive breastfeeding with employment. *Health Care for Women International* 2004;25:242-254.
211. Cattaneo A, Buzzetti R. Effect on rates of breastfeeding of training for the Baby Friendly Hospital Initiative. *BMJ* 2001; 323: 1358-1362.
212. Moran VH, Bramwell R, Dykes F, Dinwoodie K. An evaluation of skills acquisition on the WHO/UNICEF Breastfeeding Management Course using the pre-validated Breastfeeding Support Skills Tool (BeSST). *Midwifery* 2000; 16: 197-203.
213. Labarere J, Gelbert-Baudino N, Ayril AS et al. Efficacy of breastfeeding support: provided by trained clinicians during an early, routine, preventative visit: a prospective, randomized, open trial of 226 mother-infant pairs. *Pediatrics* 2005; 115: e139-146.
214. Vittoz JP, Labarere J, Castell M, Durand M, Pons JC. Effect of a training program for maternity ward professionals on duration of breastfeeding. *Birth* 2004; 31: 302-307.
215. Haggerty PA, Rutstein SO 1999. Breastfeeding and complementary infant feeding, and the postpartum effects of breastfeeding. *DHS Comparative Studies No.30*. Calverton, Maryland: Macro International Inc.
216. Mahgoup SE, Bandeke T, Nnyepi M. Breastfeeding in Botswana: practices, attitudes, patterns, and the socio-cultural factors affecting them. *J Trop Pediatr*. 2002 Aug; 48(4):195-9.
217. DeZoysa I, Rea M, Martines J. Why promote breast-feeding in diarrhoeal disease Control programmes? *Health policy and planning* 1991; 6:371-9
218. Salariya EM, Easton PM, Carter JI. Duration of breastfeeding after early initiation and frequent feeding. *Lancet* 1978; ii 1141-3
219. Lawson K, Tullock MI. Breastfeeding duration and postnatal practices. *J Adv Nurs* 1995; 22:841-9

220. Jinadu MK, Olusi SO, Agun JI, Fabiyi AK. Childhood diarrhoea in rural Nigeria: Studies on prevalence, mortality and socioenvironmental factors. *J Diarrhoeal Dis* 1991; 9:323-7
221. Ojofeitimi EO. Mothers' awareness on benefits of breast milk and cultural taboos during lactation. *Soc Sci Med* 1981; 15:135-8.
222. Calvano NR. Mode of feeding and its effects on infant mortality and morbidity. *J Trop Pediatr* 1982;28:287-93.
223. Okolo SN, Adewunmi YB, Okonji MC. Current breastfeeding knowledge, attitude, and practices of mothers in five rural communities in the Savannah region of Nigeria. *J Trop Pediatr*. 1999 Dec;45(6):323-6.
224. Victoria CG, Smith PG, Vanghan JP, Nbre LC, Lombardi C, Teixeira AM et al. Evidence for protection by breast-feeding against infant deaths from infectious disease in Brazil. *Lancet* 1987; 2:319-321
225. Serenius F, Swailm AR, Edresse AW, Hofvandrere Y. Patterns of breastfeeding weaning in Saudi Arabia. *Acta Paediatr Scand Suppl*. 1988; 121-9
226. Dettwyler KA. Infant feeding in Mali, West Africa: variation in beliefs and practice. *Social Science and Medicine* 1986;23:651-664.
227. Davies-Adetugbo AA. Sociocultural factors and the promotion of exclusive breastfeeding in rural Yoruba communities of Osun State, Nigeria. *Social Science and Medicine* 1997; 45:113-25.