

1-1-2005

# The effect of farnesol on amino acid incorporation by a wild-type and cell-wall variant strain of *Candida albicans*

Phyllis C. Braun

Fairfield University, [pcbraun@fairfield.edu](mailto:pcbraun@fairfield.edu)

Copyright National Research Council Research Press 2005.

Published version available at:

<http://nrcresearchpress.com/doi/pdf/10.1139/w05-050>

## Peer Reviewed

---

### Repository Citation

Braun, Phyllis C., "The effect of farnesol on amino acid incorporation by a wild-type and cell-wall variant strain of *Candida albicans*" (2005). *Biology Faculty Publications*. 11.

<http://digitalcommons.fairfield.edu/biology-facultypubs/11>

### Published Citation

Braun PC. 2005. The effect of farnesol on amino acid incorporation by a wild-type and cell-wall variant strain of *Candida albicans*. *Canadian Journal of Microbiology* 51(8): 715-718

This Article is brought to you for free and open access by the Biology Department at DigitalCommons@Fairfield. It has been accepted for inclusion in Biology Faculty Publications by an authorized administrator of DigitalCommons@Fairfield. For more information, please contact [digitalcommons@fairfield.edu](mailto:digitalcommons@fairfield.edu).

1

2

3

4 **The Effect of Farnesol on Amino Acid Incorporation by Wild-Type**  
5 **and Cell-Wall Variant Strain of *Candida albicans***

6

7 **Phyllis C. Braun, PhD**

8

9

10

11

12

13

14 Address Correspondence to:

15 Phyllis C. Braun, PhD

16 Biology Department

17 Fairfield University

18 1073 North Benson Road

19 Fairfield, CT 06824

20 U.S.A.

21 [pcbraun@mail.fairfield.edu](mailto:pcbraun@mail.fairfield.edu)

22

23 **ABSTRACT**

24

25 Cell-surface hydrophobicity has been shown to be an important factor in the ability of  
26 *C. albicans* to adhere to host tissue and in determining its resistance to phagocytic  
27 killing. Farnesol was recently identified as a quorum-sensing molecule that inhibits  
28 mycelial development in this organism. In this study, *C. albicans* A9 (wild type) and  
29 V10 (cell-wall-variant strain) were used. Previous experiments have indicated that  
30 A9-WT organisms have hydrophilic cell walls, while A9-V10 exhibits hydrophobic  
31 cell surfaces when both are grown in Yeast Nitrogen Broth (without amino acids)  
32 supplemented with 2% glucose at 37° C. The effect of farnesol on the incorporation  
33 of arginine, aspartic acid, glycine, histidine, leucine, and serine was studied in wild-  
34 type and cell-wall variant strains under these conditions. It was determined that both  
35 strains incorporated significantly greater amounts of all amino acids tested, except for  
36 glycine, as the concentration of farnesol increased to 10 ug/mL. Farnesol  
37 concentrations greater than 10 ug/mL abolished the enhanced effects of amino acid  
38 transport in both strains of *C. albicans*. In addition, it was determined that farnesol-  
39 treated A9-V10 hydrophobic cell surface organisms had increased amino acid uptake  
40 compared with the hydrophilic A9-WT strain although the magnitude of the effect (%  
41 increase) varied depending on the amino acid studied.

42

43 (Key Words: *Candida albicans*, farnesol, amino acid incorporation)

44           *Candida* species are ubiquitous fungi and the most common fungal pathogens  
45 affecting humans. *C. albicans* is the primary etiologic agent of candidiasis, a disease  
46 that can vary from superficial mucosal lesions in the immunocompetent host to  
47 systemic or disseminated infection in the immunocompromised individual (Odds  
48 1988; Ahearn 1978). The cell wall of the organism is important to its success as a  
49 pathogen (Odds 1985) because it is required for growth; provides rigidity and  
50 protection against osmotic insult; and is the site of contact between the organism and  
51 its surroundings. Survival of *C. albicans* in its environment is dependent on the  
52 transport of nutrients into the organism. For molecules to enter, they must first pass  
53 through the cell wall and then the plasma membrane, a limiting structure between the  
54 cell wall and the cytoplasm that acts as a permeability barrier to many solutes.

55

56           The molecular complexity of the *C. albicans* cell wall is well known. The cell  
57 wall is a heterogeneous structure composed predominantly of polysaccharides  
58 (mainly mannan, glucan, and chitin), proteins, and lipids (Calderone and Braun 1991;  
59 Odds 1985). The proteins located on the external surface of *C. albicans* are composed  
60 of high-molecular-weight mannoproteins, and constitute the outer fibrillar layer of  
61 the cell wall. These mannoproteins are regarded as important sites that influence  
62 amino acid transport by this organism (Braun 1999). Investigators have observed that  
63 the presence of fibrillar mannoproteins determines the hydrophilic nature of *C.*  
64 *albicans* cells (Glee et al. 1995; Hazen and Hazen 1992). Any changes in the length  
65 or concentration of mannoprotein fibrils, or decrease in the amount of  
66 phosphodiester-linked, acid-labile mannosyl-groups results in a hydrophobic cell  
67 surface (Masuoka and Hazen 1997).

68

69 Farnesol ( $C_{15}H_{26}O$ ), a recently discovered extracellular quorum-sensing  
70 molecule produced by *C. albicans*, blocks yeast-to-mycelium conversion when it  
71 accumulates above threshold levels. Farnesol is a hydrophobic molecule that alters  
72 cell morphology but does not alter growth rate. It is produced at all growth  
73 temperatures, and is produced continuously during growth of *C. albicans* (Hornby et  
74 al. 2001). Inhibition of *C. albicans* biofilm formation by farnesol also has been  
75 reported (Ramage et al. 2002).

76

77 The purpose of this study was to determine how amino acid incorporation was  
78 influenced by exogenous supplements of farnesol. Any changes in nutrient transport  
79 due to farnesol presence would affect the growth and presumably the pathogenesis of  
80 this organism. Two strains of *C. albicans* were used. When grown in amino-acid-  
81 deficient Yeast Nitrogen Broth (YNB) supplemented with 2% glucose, the wild-type  
82 strain of *C. albicans* (A9-WT) exhibits hydrophobic cell surfaces at 23°C and the  
83 typical hydrophilic morphology at 37°C. The cell wall variant (A9-V10) however,  
84 displays cell-surface hydrophobicity under all growth conditions (Masuoka and  
85 Hazen 1997).

86

87 *C. albicans* A9-WT and A9-V10 were maintained on brain-heart infusion  
88 slants at 4°C. The two strains were grown to mid-stationary phase in YNB buffered  
89 with 0.05-M sodium phosphate (pH 7.2) and supplemented with 2% (w/v) glucose  
90 (YNB+G) at either 23°C or 37°C to induce hydrophobic or hydrophilic cell wall  
91 conditions. Cells were washed and resuspended in PB (sodium phosphate buffer, 0.05  
92 M, pH 7.2) and kept on ice.

93

94 Cell-surface hydrophobicity (CSH) of the culture was determined by using the  
95 polystyrene microsphere assay (Hazen and Hazen 1988). A working 10% solution  
96 was prepared by placing 6 mL of stock microspheres (diameter, 0.825  $\mu\text{m}$ ; Sigma  
97 Chemical Co.) into 2.0 mL of PB, and kept on ice. The assay was performed by  
98 mixing equal volumes (100  $\mu\text{L}$ ) of washed yeast cells ( $5 \times 10^7$  cells per mL) and  
99 microspheres in polypropylene test tubes and incubating the mixture for 2 min at  
100 room temperature. The mixtures were vortexed subsequently for 30 s, and microscope  
101 slides were prepared. Cultures were determined to exhibit CSH when 95% of the  
102 population had more than three attached microspheres.

103

104 Yeast cells were resuspended in 1 mL of sterile YNB+G at a concentration of  
105  $5 \times 10^7$  cells per mL. Triplicate samples were incubated with 0, 1, or 10 micrograms  
106 of free farnesol (Sigma; trans, trans-Farnesol) for 10 minutes at 37°C. Subsequently,  
107 [ $^3\text{H}$ ] L-amino acids ([2,3- $^3\text{H}$ ]arginine, specific activity 36.8 Ci/mmol; [2,3-  
108  $^3\text{H}$ ]aspartic acid, specific activity 15.5 Ci mmol $^{-1}$ ; [2- $^3\text{H}$ ]glycine, specific activity  
109 43.8 Ci mmol $^{-1}$ ; [2,5- $^3\text{H}$ ]histidine, specific activity 50.4 Ci mmol $^{-1}$ ; [4,5- $^3\text{H}$ ]leucine,  
110 specific activity 52 Ci mmol $^{-1}$ ; [3- $^3\text{H}$ ]serine, specific activity 21.7 Ci mmol $^{-1}$ ; New  
111 England Nuclear), were individually assayed by adding the radioactively labelled  
112 substance, 1  $\mu\text{Ci ml}^{-1}$  to the cultures. These cultures were further incubated for 1 h at  
113 37°C (100 rpm). Incorporation of radiolabel was stopped by the immediate addition  
114 of 1 mL of 15% TCA. The suspensions were filtered through Whatman GF/A glass  
115 fiber filters, and the precipitate remaining on each filter was washed twice with 15%  
116 TCA and once with 95% ethanol. The filters were dried and transferred to  
117 scintillation vials, and the radioactivity was counted in ScintiSafe-LSC Cocktail.

118

119 Dry weights were determined by filtering 1-mL ( $5 \times 10^7$  cells) samples on  
120 pretared Whatman filters. These filters were washed with distilled water, dried, and  
121 weighed. Uptake of the radiolabeled nutrient is expressed as the number of picomoles  
122 incorporated per milligram of dry weight.

123

124 Statistical variables used to analyze the data included mean value, standard  
125 error of the mean, and two-tailed Student's t-test results. When two groups were  
126 compared, a *P*-value of less than 0.05 was considered to indicate a significant  
127 difference between the groups.

128

129 To determine whether A9-WT and A9-V10 strains had different growth  
130 patterns in YNB, 24-hour growth curves were performed. Calculations of the mean  
131 growth rate constant (*k*) indicated insignificant differences between wild-type and  
132 cell-wall variant at 23°C or 37°C, respectively. The mean generation time (*g*) also  
133 indicated very little growth difference between these strains at the experimental  
134 temperatures. There was, however, a significant difference in the growth rate constant  
135 and generation time for each strain at the two different temperatures (**Table 1**).

136

137 Incubation at 37°C of stationary-phase, farnesol-untreated cultures with the  
138 [<sup>3</sup>H] amino acid demonstrated that A9-V10, possessing CSH, preferentially  
139 incorporated more of the individually tested nutrient than A9-WT with a hydrophilic  
140 surface (**Figures 1 and 2**). Amino acid incorporation was linear in both strains up to  
141 60 minutes (data not shown). However, the percentage of increase varied according  
142 to the amino acid, with A9-V10 incorporating 5% more aspartic acid and leucine,  
143 11% more arginine, and 30% more histidine and serine than A9-WT. Experimental

144 results indicated A9-WT and A9-V10 cultures were unable to transport glycine under  
145 the described conditions (data not shown). Incorporation of arginine, histidine,  
146 leucine, and serine by both A9-WT and A9-V10 was further enhanced by the  
147 presence of 1 or 10 ug of farnesol compared to untreated. However, the magnitude of  
148 the effect of farnesol on transport by A9-WT and A9-V10 was dependent on the  
149 amino acid studied (**Table 2**). Aspartic acid incorporation also was enhanced by  
150 farnesol treatment; however, 10  $\mu$ g of farnesol increased A9-WT incorporation to 1.1  
151 picomoles per mg dry weight, and A9-V10 incorporation to 1.8 picomoles per mg dry  
152 weight from the untreated baseline of 0.76 and 0.8 picomoles per mg dry weight,  
153 respectively. Farnesol concentrations greater than 10  $\mu$ g/mL reduced the  
154 incorporation and presumably the transport of all amino acids tested. Increased  
155 farnesol concentrations (up to 100  $\mu$ g) abolished the augmented effects of the lower  
156 concentrations. Amino acid incorporation at these higher concentrations was reduced  
157 to base-line (0  $\mu$ g farnesol) in both strains of *C. albicans* (data not shown).

158

159

160 The role of the microenvironment apparently has a very important role in  
161 determining the pathogenicity of *C. albicans*. Farnesol, which is produced in situ by  
162 *C. albicans*, affects nutrient transport. This discovery may be relevant during in vivo  
163 infections. The results of these experiments indicate that very limited micromolar  
164 amounts of farnesol increased amino acid incorporation by this organism. The exact  
165 nature of the interaction of farnesol with *C. albicans* is not understood; however, the  
166 hydrophobic A9-V10 appears to exhibit greater nutrient transport than the A9-WT  
167 strain. This is likely a result of the lipophilic properties of farnesol, which probably  
168 enables more effective binding of this compound to a receptor site on the cell



169 membrane. Presumably, an increase in nutrient transport affects the growth,  
170 metabolism, and virulence of *C. albicans* and the morphological transformation from  
171 yeast to mycelia (a crucial step in the pathogenesis of *C. albicans*).

172         This study demonstrates that as farnesol concentrations increased, the initial  
173 incorporation enhancement was diminished to control levels (0 µg farnesol). These  
174 results indicate a gradient effect produced by farnesol that influences metabolic  
175 activities of this organism. These results, as well as previous investigations that have  
176 shown that farnesol prevents biofilm formation (Ramage et al. 2002) and inhibits  
177 mycelial development in a quorum-sensing manner (Hornby et al. 2001) suggests that  
178 farnesol might be the first of a new class of antifungal compounds.

179         In addition, Hornby et al. (2003) recently reported that compounds that block  
180 sterol biosynthesis in *C. albicans* caused an increase in intracellular and extracellular  
181 farnesol levels. Farnesol would not kill this organism and upset the balance of the  
182 normal flora in an individual. Instead, the metabolism of *C. albicans* would be  
183 affected which ultimately controls morphogenesis. Farnesol would limit *C. albicans*  
184 growth to the yeast form and in so doing restrict pathogenesis in its host. Further  
185 studies on the effect of farnesol on *C. albicans* are needed to strengthen its role as a  
186 therapeutic agent.

187

188

189

## REFERENCES

- Ahearn, D.G. 1978. Medically important yeasts. *Annu. Rev. Microbiol.* **32**:59-68.
- Braun, P.C. 1999. Nutrient uptake by *Candida albicans*: the influence of cell surface mannoproteins. *Can. J. Microbiol.* **45**:353-359.
- Calderone, R.A., Braun, P.C. 1991. Adherence and receptor relationships of *Candida albicans*. *Microbiol. Rev.* **55**:1-20.
- Glee, P.M., Sundstrom, P., Hazen, K.C. 1995. Expression of surface hydrophobic proteins by *Candida albicans* in vivo. *Infect. Immun.* **63**:1373-1379.
- Hazen, B.W. and Hazen, K.C. 1988. Dynamic expression of cell surface hydrophobicity during initial yeast cell growth and before germ tube formation of *Candida albicans*. *Infect. Immun.* **56**:2521-2525.
- Hazen, K.C. and Hazen, B.W. 1992. Hydrophobic surface protein masking by the opportunistic fungal pathogen *Candida albicans*. *Infect. Immun.* **60**:1499-1508.
- Hornby, J.M., Jensen, E.C., Lisec, A.D., Testo, J.J., Jahnke, B., Shoemaker, R., Dussault, P., Nickerson, K.W. 2001. Quorum sensing in the dimorphic fungus *Candida albicans* is mediated by farnesol. *Appl. Environ. Microbiol.* **67**:2982-2992.

Hornby, J.M., Kebaara, B.W., Nickerson, K.W. 2003. Farnesol biosynthesis in *C. albicans*: Cellular response to sterol inhibition by zaragozic acid B. *Antimicrob. Agents Chemother.* **47**:2366-2369.

Masuoka, J., Hazen, K.C. 1997. Cell wall protein mannosylation determines *Candida albicans* cell surface hydrophobicity. *Microbiology.***143**:3015-3021.

Odds, F.C. 1988. *Candida* and Candidosis. W.B. Sanders, London.

Odds, F.C. 1985. Morphogenesis in *Candida albicans*. *Crit. Rev. Microbiol.***12**:45-93.

Ramage, G., Saville, S.P., Wickes, B.L., Lopez-Ribot, J.L. 2002. Inhibition of *Candida albicans* biofilm formation by farnesol, a quorum-sensing molecule. *Appl. Environ. Microbiol.* **68**:5459-5463.

## **Acknowledgement**

P.C.B. is grateful for the lively discussions and editorial assistance of John Zoidis and the excellent technical assistance of Elizabeth Vancza and Lisa Runco.

**Table 1. Mean Growth Rate Constant (k) and Mean Generation Time (g) for *C. albicans* Strains A9-WT and A9-V10 Grown at 23°C and 37°C.**

	<u>A9-WT</u>		<u>A9-V10</u>	
	23°C	37°C	23°C	37°C
k	0.3479	0.7262	0.4592	0.7573
g (hours)	2.874	1.377	2.177	1.320


A9-WT and A9-V10 cultures were grown for 24 hours in YNB plus 2% glucose broth. Values represent the average of 5 experimental growth curves. Data were analyzed using Student's t-test. When growth comparisons were determined, a *P*-value less than 0.05 was considered to indicate a significant difference between the groups.

**Table 2. Magnitude of Effect (% Increase) of Farnesol 1  $\mu$ g and 10  $\mu$ g on Amino Acid Transport by *C. albicans* Strains A9-WT and A9-V10.**


	<b>A9-WT</b>		<b>A9-V10</b>	
	<b>Farnesol 1 <math>\mu</math>g</b>	<b>Farnesol 10 <math>\mu</math>g</b>	<b>Farnesol 1 <math>\mu</math>g</b>	<b>Farnesol 10 <math>\mu</math>g</b>
Arginine	2	26	4	19
Histidine	14	23	10	21
Leucine	21	42	22	55
Serine	44	54	33	90

**Figure 1**

## Arginine




# Histidine




**Figure 2**



# Leucine



# Serine



## Legend

**Figs. 1-2:**

Incorporation of four different L-[<sup>3</sup>H] amino acids by pretreated farnesol *C. albicans* strains A9-WT(hydrophilic) and A9-V10 (hydrophobic) was determined after a 1 h incubation at 37°C. Values represent the SEM of triplicate samples from several experiments (n=6). Student's *t* test was used to analyze the data. When incorporation comparisons between cell strains for a particular amino acid was determined, a  $P \leq 0.05$  was considered to indicate a significant difference between groups.