

1-1-2004

Observation of an exotic baryon with $S = +1$ in photoproduction from the proton

V. Kubarovsky

Angela Biselli

Fairfield University, abiselli@fairfield.edu

CLAS Collaboration

Copyright American Physical Society Publisher final version available at <http://prl.aps.org/pdf/PRL/v92/i3/e032001> and <http://journals.aps.org/prl/pdf/10.1103/PhysRevLett.92.049902>

Peer Reviewed

Repository Citation

Kubarovsky, V.; Biselli, Angela; and CLAS Collaboration, "Observation of an exotic baryon with $S = +1$ in photoproduction from the proton" (2004). *Physics Faculty Publications*. 91.
<http://digitalcommons.fairfield.edu/physics-facultypubs/91>

Published Citation

V. Kubarovsky et al. [CLAS Collaboration], "Observation of an exotic baryon with $S = +1$ in photoproduction from the proton", *Phys. Rev. Lett.* 92, 032001 (2004) [Erratum- *ibid.* 92, 049902 (2004)] DOI: 10.1103/PhysRevLett.92.032001

This Article is brought to you for free and open access by the Physics Department at DigitalCommons@Fairfield. It has been accepted for inclusion in Physics Faculty Publications by an authorized administrator of DigitalCommons@Fairfield. For more information, please contact digitalcommons@fairfield.edu.

Observation of an Exotic Baryon with $S = +1$ in Photoproduction from the Proton

V. Kubarovsky,^{1,3} L. Guo,² D. P. Weygand,³ P. Stoler,¹ M. Battaglieri,¹⁸ R. DeVita,¹⁸ G. Adams,¹ Ji Li,¹ M. Nozar,³ C. Salgado,²⁶ P. Ambrozewicz,¹³ E. Anciant,⁵ M. Anghinolfi,¹⁸ B. Asavapibhop,²⁴ G. Audit,⁵ T. Auger,⁵ H. Avakian,³ H. Bagdasaryan,²⁸ J. P. Ball,⁴ S. Barrow,¹⁴ K. Beard,²¹ M. Bektasoglu,²⁷ M. Bellis,¹ N. Benmouna,¹⁵ B. L. Berman,¹⁵ N. Bianchi,¹⁷ A. S. Biselli,⁷ S. Boiarinov,²⁰ S. Bouchigny,¹⁹ R. Bradford,⁷ D. Branford,¹² W. J. Briscoe,¹⁵ W. K. Brooks,³ V. D. Burkert,³ C. Butuceanu,³⁷ J. R. Calarco,²⁵ D. S. Carman,⁷ B. Carnahan,⁸ C. Cetina,¹⁵ S. Chen,¹⁴ L. Ciciani,²⁸ P. L. Cole,³³ J. Connelly,¹⁵ D. Cords,^{3,*} P. Corvisiero,¹⁸ D. Crabb,³⁶ H. Crannell,⁸ J. P. Cummings,¹ E. De Sanctis,¹⁷ P. V. Degtyarenko,³ H. Denizli,²⁹ L. Dennis,¹⁴ K. V. Dharmawardane,²⁸ C. Djalali,³² G. E. Dodge,²⁸ D. Doughty,⁹ P. Dragovitsch,¹⁴ M. Dugger,⁴ S. Dytman,²⁹ O. P. Dzyubak,³² H. Egiyan,³ K. S. Egiyan,³⁸ L. Elouadrhiri,⁹ A. Empl,¹ P. Eugenio,¹⁴ L. Farhi,⁵ R. Fatemi,³⁶ R. J. Feuerbach,⁷ J. Ficencic,³⁵ T. A. Forest,²⁸ V. Frolov,¹ H. Funsten,³⁷ S. J. Gaff,¹¹ M. Garçon,⁵ G. Gavalian,²⁵ G. P. Gilfoyle,³¹ K. L. Giovanetti,²¹ P. Girard,³² R. Gothe,³² C. I. O. Gordon,¹⁶ K. Griffioen,³⁷ M. Guidal,¹⁹ M. Guillo,³² V. Gyurjyan,³ C. Hadjidakis,¹⁹ R. S. Hakobyan,⁸ D. Hancock,³⁷ J. Hardie,⁹ D. Heddle,⁹ P. Heimberg,¹⁵ F. W. Hersman,²⁵ K. Hicks,²⁷ M. Holtrop,²⁵ J. Hu,¹ Y. Ilieva,¹⁵ M. M. Ito,³ D. Jenkins,³⁵ K. Joo,¹⁰ H. G. Juengst,¹⁵ J. H. Kelley,¹¹ M. Khandaker,²⁶ K. Y. Kim,²⁹ K. Kim,²² W. Kim,²² F. J. Klein,³ A. V. Klimenko,²⁸ M. Klusman,¹ M. Kossov,²⁰ L. H. Kramer,¹³ S. E. Kuhn,²⁸ J. Kuhn,⁷ J. Lachniet,⁷ J. M. Laget,⁵ J. Langheinrich,³² D. Lawrence,²⁴ A. Longhi,⁸ K. Lukashin,³ R. W. Major,^{31,*} J. J. Manak,³ C. Marchand,⁵ S. McAleer,¹⁴ J. W. C. McNabb,⁷ B. A. Mecking,³ S. Mehrabyan,²⁹ J. J. Melone,¹⁶ M. D. Mestayer,³ C. A. Meyer,⁷ K. Mikhailov,²⁰ R. Minehart,³⁶ M. Mirazita,¹⁷ R. Miskimen,²⁴ V. Mokeev,³⁹ L. Morand,⁵ S. A. Morrow,⁵ M. U. Mozer,²⁷ V. Muccifora,¹⁷ J. Mueller,²⁹ G. S. Mutchler,³⁰ J. Napolitano,¹ R. Nasseripour,¹³ S. O. Nelson,¹¹ S. Niccolai,¹⁵ G. Niculescu,²⁷ I. Niculescu,²¹ B. B. Niczyporuk,³ R. A. Niyazov,²⁸ J. T. O'Brien,⁸ G. V. O'Rielly,¹⁵ A. K. Opper,²⁷ M. Osipenko,¹⁸ K. Park,²² E. Pasyuk,⁴ G. Peterson,²⁴ S. A. Philips,¹⁵ N. Pivnyuk,²⁰ D. Pocanic,³⁶ O. Pogorelko,²⁰ E. Polli,¹⁷ S. Pozdniakov,²⁰ B. M. Preedom,³² J. W. Price,⁶ Y. Prok,³⁶ D. Protopopescu,¹⁶ L. M. Qin,²⁸ B. A. Raue,¹³ G. Riccardi,¹⁴ M. Ripani,¹⁸ B. G. Ritchie,⁴ F. Ronchetti,¹⁷ P. Rossi,¹⁷ D. Rowntree,²³ P. D. Rubin,³¹ F. Sabatié,⁵ K. Sabourov,¹¹ J. P. Santoro,³⁵ V. Sapunenko,¹⁸ M. Sargsyan,¹³ R. A. Schumacher,⁷ V. S. Serov,²⁰ A. Shafi,¹⁵ Y. G. Sharabian,³⁸ J. Shaw,²⁴ S. Simionatto,¹⁵ A. V. Skabelin,²³ E. S. Smith,³ T. Smith,²⁵ L. C. Smith,³⁶ D. I. Sober,⁸ M. Spraker,¹¹ A. Stavinsky,²⁰ S. Stepanyan,³⁸ I. I. Strakovsky,¹⁵ S. Strauch,¹⁵ M. Taiuti,¹⁸ S. Taylor,³⁰ D. J. Tedeschi,³² U. Thoma,³ R. Thompson,²⁹ L. Todor,⁷ C. Tur,³² M. Ungaro,¹ M. F. Vineyard,³⁴ A. V. Vlassov,²⁰ K. Wang,³⁶ L. B. Weinstein,²⁸ A. Weisberg,²⁷ C. S. Whisnant,³² E. Wolin,³ M. H. Wood,³² A. Yegneswaran,³ and J. Yun²⁸

(CLAS Collaboration)

¹Rensselaer Polytechnic Institute, Troy, New York 12180-3590, USA

²Vanderbilt University, Nashville, Tennessee 37235, USA

³Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606, USA

⁴Arizona State University, Tempe, Arizona 85287-1504, USA

⁵CEA-Saclay, Service de Physique Nucléaire, F91191 Gif-sur-Yvette, Cedex, France

⁶University of California at Los Angeles, Los Angeles, California 90095-1547, USA

⁷Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA

⁸Catholic University of America, Washington, D.C. 20064, USA

⁹Christopher Newport University, Newport News, Virginia 23606, USA

¹⁰University of Connecticut, Storrs, Connecticut 06269, USA

¹¹Duke University, Durham, North Carolina 27708-0305, USA

¹²Edinburgh University, Edinburgh EH9 3JZ, United Kingdom

¹³Florida International University, Miami, Florida 33199, USA

¹⁴Florida State University, Tallahassee, Florida 32306, USA

¹⁵The George Washington University, Washington, D.C. 20052, USA

¹⁶University of Glasgow, Glasgow G12 8QQ, United Kingdom

¹⁷INFN, Laboratori Nazionali di Frascati, Frascati, Italy

¹⁸INFN, Sezione di Genova, 16146 Genova, Italy

¹⁹Institut de Physique Nucleaire ORSAY, Orsay, France

²⁰Institute of Theoretical and Experimental Physics, Moscow, 117259, Russia

²¹James Madison University, Harrisonburg, Virginia 22807, USA

²²Kyungpook National University, Daegu 702-701, South Korea

²³*Massachusetts Institute of Technology, Cambridge, Massachusetts 02139-4307, USA*²⁴*University of Massachusetts, Amherst, Massachusetts 01003, USA*²⁵*University of New Hampshire, Durham, New Hampshire 03824-3568, USA*²⁶*Norfolk State University, Norfolk, Virginia 23504, USA*²⁷*Ohio University, Athens, Ohio 45701, USA*²⁸*Old Dominion University, Norfolk, Virginia 23529, USA*²⁹*University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA*³⁰*Rice University, Houston, Texas 77005-1892, USA*³¹*University of Richmond, Richmond, Virginia 23173, USA*³²*University of South Carolina, Columbia, South Carolina 29208, USA*³³*University of Texas at El Paso, El Paso, Texas 79968, USA*³⁴*Union College, Schenectady, New York 12308, USA*³⁵*Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061-0435, USA*³⁶*University of Virginia, Charlottesville, Virginia 22901, USA*³⁷*College of William and Mary, Williamsburg, Virginia 23187-8795, USA*³⁸*Yerevan Physics Institute, 375036 Yerevan, Armenia*³⁹*General Nuclear Physics Institute, Moscow State University, 119899 Moscow, Russia*

(Received 21 November 2003; published 21 January 2004; publisher error corrected 22 January 2004)

The reaction $\gamma p \rightarrow \pi^+ K^- K^+ n$ was studied at Jefferson Laboratory using a tagged photon beam with an energy range of 3–5.47 GeV. A narrow baryon state with strangeness $S = +1$ and mass $M = 1555 \pm 10 \text{ MeV}/c^2$ was observed in the nK^+ invariant mass spectrum. The peak's width is consistent with the CLAS resolution (FWHM = 26 MeV/ c^2), and its statistical significance is $(7.8 \pm 1.0)\sigma$. A baryon with positive strangeness has exotic structure and cannot be described in the framework of the naive constituent quark model. The mass of the observed state is consistent with the mass predicted by the chiral soliton model for the Θ^+ baryon. In addition, the pK^+ invariant mass distribution was analyzed in the reaction $\gamma p \rightarrow K^- K^+ p$ with high statistics in search of doubly charged exotic baryon states. No resonance structures were found in this spectrum.

DOI: 10.1103/PhysRevLett.92.032001

PACS numbers: 13.60.Rj, 14.20.Jn, 14.80.-j

The constituent quark model describes light mesons as bound states of a quark and an antiquark ($q\bar{q}$), and baryons as bound states of three quarks (qqq), where q is u , d , or s . It is a remarkable feature of meson and baryon spectroscopy that practically all well-established particles can be categorized using this naive model. At the same time, quantum chromodynamics (QCD) predicts the existence of so-called exotic mesons and baryons with more complicated internal structures. Exotic mesons may be classified such as glueballs (ggg), hybrids ($q\bar{q}g$), and four-quark ($q\bar{q}q\bar{q}$) states, and exotic baryons as ($qqqg$) or ($qqqq\bar{q}$) states.

A baryon with strangeness quantum number $S = +1$ is an excellent example of a particle whose exotic structure is manifest. Diakonov, Petrov, and Polyakov [1], in the framework of the chiral soliton model, have predicted a spin 1/2, isospin 0, and strangeness $S = +1$ exotic baryon Θ^+ with mass $M \sim 1.53 \text{ GeV}/c^2$. Also, the pentaquark state was considered in the quark models [2] and lattice QCD [3]. The possible quark structure of Θ^+ is ($uudd\bar{s}$).

Experimental evidence for a narrow $S = +1$ baryon state has been reported in the interactions of photons and kaons with nuclear targets: $\gamma n \rightarrow K^+ K^- n$ (on a ^{12}C target) [4], $K^+ \text{Xe}$ collisions in the pK^0 decay mode [5], and the exclusive photodeuteron interaction in the nK^+ decay mode [6]. In all these reactions the primary interaction is with a neutron in the initial state. The CLAS

Collaboration has reported the observation of the same state in the photoproduction reaction from the proton target [7]. The SAPHIR Collaboration [8] has also reported the observation of this state in the reaction $\gamma p \rightarrow nK^+ K_s^0$. A narrow peak with mass around $M = 1.54 \text{ GeV}/c^2$ and width less than $25 \text{ MeV}/c^2$ was observed and interpreted as the production and decay of the exotic Θ^+ baryon. More recently, the ITEP Collaboration observed a narrow baryon resonance in the invariant mass of the pK_s^0 system formed in neutrino and antineutrino collisions with nuclei [9].

This Letter reports a more comprehensive study than in [7] of the Θ^+ production on a proton target which includes data from three distinct runs under different experimental conditions in CLAS [10]. Two reactions, $\gamma p \rightarrow \pi^+ K^+ K^- n$ and $\gamma p \rightarrow K^+ K^- p$, have been analyzed. Of the three runs, run (a) and run (b) had identical geometrical acceptance and trigger requirement with run (c) having slightly different running conditions. Runs (a), (b), and (c) had a tagged photon beam in the energy range of 3.2–3.95, 3–5.25, and 4.8–5.47 GeV, respectively [11]. CLAS is a sixfold segmented toroidal magnetic spectrometer (details described in [10]). Run (c) triggered on the events with at least two out of six CLAS sectors having charged tracks, while runs (a) and (b) triggered on events with hits in opposite sectors. Runs (a) and (b) had the hydrogen target in the standard position, but in run (c) the target was moved upstream by




FIG. 1 (color online). The missing mass distributions in the reaction $\gamma p \rightarrow \pi^+ K^- K^+ X$ for the three different runs [runs (a), (b), and (c)] and the combined data spectrum [(d)]. The mass resolutions are 10, 11, and 19 MeV/c^2 for runs (a), (b), and (c), respectively.

1 m to improve the CLAS acceptance in the forward direction, especially for the negative charged particles. The estimated integrated luminosity for the combined data of runs (a) and (b) is 2 pb^{-1} , and run (c) is 2.7 pb^{-1} . The combined analysis of these three runs offers access to a wider range of acceptance and energies.

Events having a π^+ , K^+ , and K^- in the final state, identified by time of flight, were selected for the analysis of the reaction $\gamma p \rightarrow \pi^+ K^- K^+ n$. The missing mass distributions for the reaction $\gamma p \rightarrow \pi^+ K^- K^+ X$ are shown in Fig. 1. A neutron peak is clearly seen in each of these distributions. The fitted masses of the neutron peaks are consistent with each other within $3 \text{ MeV}/c^2$. The mass resolution for runs (a) and (b) is about twice as good as in run (c) due to the fact that run (c) had higher energy and the magnetic field was reduced by a factor of 2. Events within $\pm 2\sigma$ of the neutron peak were retained for each run individually, resulting in a total of 14 000 events.

There are about 200 ϕ mesons in the selected sample, which were removed by eliminating events with the $K^+ K^-$ effective mass less than $1.06 \text{ GeV}/c^2$. The final nK^+ invariant mass spectrum calculated from missing mass in the reaction $\gamma p \rightarrow \pi^+ K^- X$, combining data from all three data runs, is shown in Fig. 2. No obvious structure is seen in this spectrum.

We explored various possible t -channel processes to understand the potential production mechanisms for the Θ^+ as well as the background, examples of which are illustrated in Fig. 3. A peak appears most clearly when requiring $\cos\theta_{\pi^+}^* > 0.8$, where $\theta_{\pi^+}^*$ is the center-of-mass angle between the π^+ and the photon beam. This requirement approximately corresponds to $-t < 0.28 \text{ GeV}/c^2$ where $t = (k - p)^2$, k is the photon 4-momentum, and p




FIG. 2. The nK^+ invariant mass spectrum in the reaction $\gamma p \rightarrow \pi^+ K^- K^+ (n)$. The neutron was measured from the missing 4-momentum.

is the pion 4-momentum. This would correspond to an enhancement of the t -channel process as shown in Fig. 3(a). The spectrum is shown in the inset in Fig. 4. As a systematic check, we varied the pion angular cut from 0.7 to 0.9 and found that in all cases the peak was clearly visible.

The background reaction $\gamma p \rightarrow \pi^+ K^- K^+ n$ is dominated by meson resonance production decaying to $K^+ K^-$ with the excitation of baryon resonances decaying to $n\pi^+$, or meson resonance production decaying to $K^+ K^- \pi^+$, both with small momentum transfer to the




FIG. 3 (color online). The diagram in (a) shows a possible production mechanism for the Θ^+ , which could be a decay product of an intermediate baryon resonance. The three diagrams in (b)–(d) are background processes in the reaction $\gamma p \rightarrow \pi^+ K^- K^+ (n)$. All background processes have a K^+ going in the forward direction in the center-of-mass system.




FIG. 4 (color online). The nK^+ invariant mass spectrum in the reaction $\gamma p \rightarrow \pi^+ K^- K^+(n)$ with the cut $\cos\theta_{\pi^+}^* > 0.8$ and $\cos\theta_{K^+}^* < 0.6$. $\theta_{\pi^+}^*$ and $\theta_{K^+}^*$ are the angles between the π^+ and K^+ mesons and photon beam in the center-of-mass system. The background function we used in the fit was obtained from the simulation. The inset shows the nK^+ invariant mass spectrum with only the $\cos\theta_{\pi^+}^* > 0.8$ cut.

meson system. These processes have the K^+ moving forward in the center-of-mass system [Figs. 3(b)–3(d)]. To suppress such backgrounds, a cut was applied to eliminate events having a positive kaon going forward with $\cos\theta_{K^+}^* > 0.6$, where $\theta_{K^+}^*$ is the center-of-mass angle between the K^+ and the photon beam. The remaining data sample is virtually free of the contaminating events that have baryons decaying to $n\pi^+$ in this final state since the π^+ from such an event will most likely not move very forward in the center-of-mass system. The Θ^+ peak was clearly observed in each of the three data sets; the resulting nK^+ mass spectra are combined and shown in Fig. 4.

An investigation has been conducted to test whether a narrow peak in the nK^+ invariant mass spectrum can be artificially manufactured. First, we checked the sidebands around the neutron in Fig. 1; the resulting nK^+ effective mass distribution is structureless. We also considered the effect of the kinematic requirements that we applied by performing a Monte Carlo simulation based on $nK^+K^-\pi^+$ 4-body phase space, $nK^+\bar{K}_0^*$ 3-body phase space, and t -channel meson production. The meson events in the latter process are generated using $K^+K^-\pi^+$ 3-body phase space and the shape of the $K^+K^-\pi^+$ invariant mass distribution from the data. We found no structure generated using the same cuts on the simulated events as applied on the data. Furthermore, one may consider whether some particular combination of meson waves can be reflected as a narrow peak in the nK^+ invariant mass distribution. We performed a full partial wave analysis of the $K^+K^-\pi^+$ meson system on run (c) data and utilized the prediction of this analysis to further probe the possibility of meson reflection into the nK^+

invariant mass spectrum. Again, we found that with a set of meson partial waves that well describes the entire data set we did not generate a narrow Θ^+ peak when the same angular cuts as above are applied.

The final nK^+ effective mass distribution (Fig. 4) was fitted by the sum of a Gaussian function and a background function obtained from the simulation. The fit parameters are $N_{\Theta^+} = 41 \pm 10$, $M = 1555 \pm 1 \text{ MeV}/c^2$, and $\Gamma = 26 \pm 7 \text{ MeV}/c^2$ (FWHM), where the errors are statistical. The systematic mass scale uncertainty is estimated to be $\pm 10 \text{ MeV}/c^2$. This uncertainty is larger than our previously reported uncertainty [6] because of the different energy range and running conditions and is mainly due to the momentum calibration of the CLAS detector and the photon beam energy calibration. The statistical significance for the fit in Fig. 4 over a $40 \text{ MeV}/c^2$ mass window is calculated as $N_P/\sqrt{N_B}$, where N_B is the number of counts in the background fit under the peak and N_P is the number of counts in the peak. We estimate the significance to be $7.8 \pm 1.0\sigma$. The uncertainty of 1.0σ is due to the different background functions that we tried. When a simple polynomial background is used, the statistical significance is higher. In the present analysis we used the background function obtained from the simulation as discussed above. The fact that the angular cuts we applied enhanced the Θ^+ signal suggests the possible production of an N^*/Δ^* that decays to Θ^+ and K^- . If the Θ^+ is an isosinglet, the intermediate state can be only an N^* . The nK^+K^- invariant mass is shown in Fig. 5 for the events with nK^+ effective mass between 1.54 and $1.58 \text{ GeV}/c^2$. The apparent excess of events near $2.4 \text{ GeV}/c^2$ is suggestive of an intermediate baryon state. A possible production mechanism that could contribute to the Θ^+ production is shown in Fig. 3(a). Similar processes were also theoretically considered in Ref. [12]. The simulation we described previously also demonstrated that the angular cuts we applied could not generate a narrow peak in the nK^+K^- invariant mass spectrum from any of the three data runs.

In addition, a search for a manifestly exotic baryon ($Q = 2$, $S = +1$) was performed in the reaction $\gamma p \rightarrow K^- X^{++}$, $X^{++} \rightarrow pK^+$. There were 225 000 events with a proton and K^+ in the final state, which were selected for the analysis of this reaction. The K^- was identified by the missing mass technique. After the removal of $\phi \rightarrow K^+K^-$ and $\Lambda(1520) \rightarrow K^-p$, we observe no resonant structures in the pK^+ invariant mass distribution for the remaining 130 000 events. The pK^+ invariant mass spectra for different $\cos\theta_{K^-}^*$ were analyzed as well, where $\theta_{K^-}^*$ is the angle between the K^- and incident photon in the center-of-mass system. There are no resonance structures evident in any of these distributions. A more detailed analysis will be presented in a future paper.

In summary, the reaction $\gamma p \rightarrow \pi^+ K^- K^+ n$ was studied at Jefferson Laboratory with photon energies from 3 to 5.47 GeV using the CLAS detector. A narrow




FIG. 5 (color online). The nK^+K^- invariant mass spectrum calculated from the missing mass off the π^+ in the reaction $\gamma p \rightarrow \pi^+ K^- K^+(n)$ with the cuts $\cos\theta_{\pi^+}^* > 0.8$ and $\cos\theta_{K^+}^* < 0.6$. $\theta_{\pi^+}^*$ and $\theta_{K^+}^*$ are the angles between the π^+ or K^+ mesons and photon beam in the center-of-mass system. These events have $M(K^+n)$ between 1.54 and 1.58 GeV/c^2 . The shape of the background curve was obtained from the simulation as discussed in the text. The inset shows the nK^+K^- invariant mass spectrum for all other events in Fig. 4.

baryon state with positive strangeness $S = +1$, mass $M = 1555 \pm 10 \text{ MeV}/c^2$, and width $\Gamma < 26 \text{ MeV}/c^2$ (FWHM) was observed. The width is close to the experimental mass resolution of the CLAS detector. The peak's statistical significance is $(7.8 \pm 1.0)\sigma$. In addition, the pK^+ invariant mass distribution was analyzed in the reaction $\gamma p \rightarrow K^- K^+ p$ with high statistics. No resonance structures were found in this spectrum.

We acknowledge the outstanding efforts of the staff of the Accelerator and the Physics Divisions at Jefferson Laboratory that made this experiment possible. This work was supported in part by the U.S. Department of Energy, by the National Science Foundation, by the Istituto Nazionale di Fisica Nucleare, by the French Centre National de la Recherche Scientifique, by the French Commissariat à l'Énergie Atomique, by an Emmy Noether grant from the Deutsche Forschungsgemeinschaft, and by the Korean Science and Engineering Foundation. The Southeastern Universities Research Association (SURA) operates the Thomas Jefferson National Accelerator Facility for the United States Department of Energy under Contract No. DE-AC05-84ER40150.

*Deceased.

- [1] D. Diakonov, V. Petrov, and M. Polyakov, *Z. Phys. A* **359**, 305 (1997).
- [2] M. Karliner and H.J. Lipkin, *Phys. Lett. B* **575**, 249 (2003); R.L. Jaffe and F. Wilczek, *Phys. Rev. Lett.* **91**, 232003 (2003).
- [3] F. Csikor *et al.*, *J. High Energy Phys.* 0311 (2003) 070.
- [4] T. Nakano *et al.*, *Phys. Rev. Lett.* **91**, 012002 (2003).
- [5] V. Barmin *et al.*, *Phys. At. Nucl.* **66**, 1715 (2003).
- [6] S. Stepanyan *et al.*, *Phys. Rev. Lett.* **91**, 252001 (2003).
- [7] V. Kubarovsky and S. Stepanyan, hep-ex/0307088.
- [8] J. Barth *et al.*, *Phys. Lett. B* **572**, 127 (2003).
- [9] A. E. Asratyan, A. G. Dolgolenko, and M. A. Kubantsev, hep-ex/0309042.
- [10] B. A. Mecking *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **503**, 513 (2003).
- [11] D. Sober *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **440**, 263 (2000).
- [12] P. Page, hep-ph/0310200.